

IMBEDDING THE IRRATIONAL ROTATION C*-ALGEBRA INTO AN AF-ALGEBRA

M. PIMSNER and D. VOICULESCU

Let $\theta \in (0, 1)$ be an irrational number and A_θ the corresponding irrational rotation C*-algebra (i.e. the cross-product of the continuous functions on the unit circle by the automorphism corresponding to the rotation of angle $2\pi\theta$). This C*-algebra is simple and has a unique trace state τ . It has been shown by M. A. Rieffel [9] that for every $\alpha \in [0, 1] \cap (\mathbf{Z} + \theta\mathbf{Z})$ there is a projection $p \in A_\theta$ such that $\tau(p) = \alpha$ and he conjectured that $\tau(p)$ can take values only in $[0, 1] \cap (\mathbf{Z} + \theta\mathbf{Z})$ and even more that the map $K_0(A_\theta) \rightarrow \mathbf{R}$ induced by τ establishes an isomorphism between the ordered group $K_0(A_\theta)$ and the subgroup $\mathbf{Z} + \theta\mathbf{Z}$ of \mathbf{R} . The group $K_0(A_\theta)$ has also appeared in the context of A. Connes' work [2] on operator algebras associated with foliations, as the range of an index-map in the case of the Kronecker flow on the 2-torus.

In the present note we construct an imbedding of A_θ into an AF-algebra constructed by E. G. Effros and C. L. Shen ([3]), the K-group of which is $\mathbf{Z} + \theta\mathbf{Z}$. This has as a consequence the fact that $\tau(p) \in \mathbf{Z} + \theta\mathbf{Z}$ for all projections $p \in A_\theta$, or somewhat stronger, that the image of the homomorphism $K_0(A_\theta) \rightarrow \mathbf{R}$ induced by τ is contained in $\mathbf{Z} + \theta\mathbf{Z}$ thus confirming a part of the facts conjectured by M. A. Rieffel. This, together with Rieffel's result, implies that A_{θ_1} and A_{θ_2} are isomorphic if and only if $\theta_1 \in \{\theta_2, 1 - \theta_2\}$.

The irrational rotation algebra is generated by two unitaries $u, v \in A_\theta$ such that $u^*vu = e^{2\pi i\theta}v$ and every pair U, V of unitaries in some unital C*-algebra B , satisfying $U^*VU = e^{2\pi i\theta}V$ determines a *-monomorphism $\psi: A_\theta \rightarrow B$, such that $\psi(v) = V, \psi(u) = U$. This fact will be used in what follows.

Let $[a_1, a_2, \dots, a_n, \dots]$ be the continued fraction expansion of θ (see [8]). The n 'th convergent $\frac{p_n}{q_n}$ equals $[a_1, a_2, \dots, a_n]$ and p_n, q_n may be computed recurrently by the formulae:

$$(1) \quad \begin{aligned} p_n &= a_n p_{n-1} + p_{n-2} & (n \geq 2) \text{ with } p_0 = 0, p_1 = 1 \\ q_n &= a_n q_{n-1} + q_{n-2} & (n \geq 2) \text{ with } q_0 = 1, q_1 = a_1. \end{aligned}$$

For each $n \in \mathbb{N}$ let \mathcal{M}_{q_n} denote the $q_n \times q_n$ matrices acting on the Hilbert space \mathbb{C}^{q_n} . The canonical basis in \mathbb{C}^{q_n} will be denoted by $e(n; j)$, $1 \leq j \leq q_n$. The AF -algebra into which we shall imbed A_θ is the inductive limit

$$(2) \quad \rightarrow A_{n-1} \xrightarrow{\rho_n} A_n \rightarrow$$

of the finite-dimensional C^* -algebras $A_n = \mathcal{M}_{q_n} \oplus \mathcal{M}_{q_{n-1}}$, the imbeddings $\rho_n: A_{n-1} \rightarrow A_n$ being defined by

$$\rho_n(x \oplus y) = W_n(x \oplus \dots \oplus x \oplus y) W_n^* \oplus x$$

a_n -times

where $x \in \mathcal{M}_{q_{n-1}}$, $y \in \mathcal{M}_{q_{n-2}}$ and

$$W_n: \underbrace{\mathbb{C}^{q_{n-1}} \oplus \dots \oplus \mathbb{C}^{q_{n-1}}}_{a_n\text{-times}} \oplus \mathbb{C}^{q_{n-2}} \rightarrow \mathbb{C}^{q_n}$$

is a unitary operator we shall define below. Remark that (1) makes the definition of W_n possible.

To define W_n we fix an integer s , satisfying $q_{n-2}/4 \leq s \leq q_{n-2}/2$. This is always possible when $n \geq 6$, so that the inductive limit (2) will start at $A_5 \rightarrow A_6 \rightarrow A_7 \dots$.

The operator W_n ($n \geq 6$) is of the form:

$$\begin{aligned} W_n(\xi_1 \oplus \dots \oplus \xi_{a_n} \oplus \eta) = \\ = W_n^{(1)}\xi_1 + \dots + W_n^{(a_n)}\xi_{a_n} + \tilde{W}_n \eta \end{aligned}$$

where

$$\begin{aligned} W_n^{(k)}: \mathbb{C}^{q_{n-1}} &\rightarrow \mathbb{C}^{q_n}, \quad (1 \leq k \leq a_n) \\ \tilde{W}_n: \mathbb{C}^{q_{n-2}} &\rightarrow \mathbb{C}^{q_n} \end{aligned}$$

are isometric operators given by the formulae:

$$\begin{aligned} W_n^{(k)}e(n-1; j) = \alpha_j^{(k)}e(n; a(k, j)) + \beta_j^{(k)}e(n; b(k, j)) + \gamma_j^{(k)}e(n; c(k, j)) \\ (1 \leq j \leq q_{n-1}, \quad 1 \leq k \leq a_n) \end{aligned}$$

and

$$\begin{aligned} \tilde{W}_n e(n-2; j) = \lambda_j e(n; l(j)) + \mu_j e(n; m(j)) \\ (1 \leq j \leq q_{n-2}) \end{aligned}$$

with indices and coefficients given by formulae listed below.

The indices are defined modulo q_n by:

$$a(k, j) = (-1)^k \left[\frac{k}{2} \right] q_{n-1} + j$$

$$b(k, j) = (-1)^{k+1} \left[\frac{k}{2} \right] q_{n-1} + j$$

$$c(k, j) = (-1)^k \left[\frac{k}{2} \right] q_{n-1} - q_{n-1} + j$$

for $1 \leq k \leq a_n, 1 \leq j \leq q_{n-1}$ and

$$l(j) = (-1)^{a_n+1} \left[\frac{a_n+1}{2} \right] q_{n-1} + j$$

$$m(j) = \left[\frac{a_n+1}{2} \right] q_{n-1} + j - \frac{(1 - (-1)^{a_n})}{2} q_{n-2}$$

for $1 \leq j \leq q_{n-2}$ and with $[x]$ denoting the integer part of x .

The formulae for the coefficients are:

$$\alpha_j^{(1)} = 0 \quad \text{where } 1 \leq j \leq q_{n-1}$$

and for $1 < k \leq a_n$

$$\alpha_j^{(k)} = \begin{cases} \exp\left(\frac{\pi i}{s} \cdot j \cdot \frac{1 - (-1)^k}{2}\right) \cos\left(\frac{\pi}{2s} j\right) & \text{when } 1 \leq j \leq s \\ 0 & \text{when } s < j \leq q_{n-1}; \end{cases}$$

$$\beta_j^{(1)} = \begin{cases} 1 & \text{when } 1 \leq j \leq q_{n-1} - s \\ \cos\left(\frac{\pi}{2s} (j - q_{n-1} + s)\right) & \text{when } q_{n-1} - s < j \leq q_{n-1} \end{cases}$$

and for $1 < k \leq a_n$

$$\beta_j^{(k)} = \begin{cases} (-1)^k \exp\left(\frac{\pi i}{s} \cdot j \cdot \frac{1 - (-1)^k}{2}\right) \sin\left(\frac{\pi}{2s} j\right) & \text{when } 1 \leq j \leq s \\ 1 & \text{when } s < j \leq q_{n-1} - s \\ \exp\left(\frac{\pi i}{s} (j - q_{n-1} + s) \frac{1 + (-1)^k}{2}\right) \cos\left(\frac{\pi}{2s} (j - q_{n-1} + s)\right) & \text{when } q_{n-1} - s < j \leq q_{n-1} \end{cases}$$

and for $1 \leq k \leq a_n$

$$y_j^{(k)} = \begin{cases} 0 & \text{when } 1 \leq j \leq q_{n-1} - s \\ (-1)^{k+1} \exp\left(\frac{\pi i}{s} (j - q_{n-1} + s) \frac{1 + (-1)^k}{2}\right) \sin\left(\frac{\pi}{2s} (j - q_{n-1} + s)\right) & \text{when } q_{n-1} - s < j \leq q_{n-1}. \end{cases}$$

The formulae for λ_j and μ_j are distinct according to the fact that a_n is even or odd.

For a_n an even number, we have:

$$\lambda_j = \begin{cases} \exp\left(\frac{\pi i}{s} j\right) \cos\left(\frac{\pi}{2s} j\right) & \text{when } 1 \leq j \leq s \\ 0 & \text{when } s < j \leq q_{n-2} \end{cases}$$

$$\mu_j = \begin{cases} -\exp\left(\frac{\pi i}{s} j\right) \sin\left(\frac{\pi}{2s} j\right) & \text{when } 1 \leq j \leq s \\ 1 & \text{when } s < j \leq q_{n-2}. \end{cases}$$

For a_n an odd number, we have:

$$\lambda_j = \begin{cases} 1 & \text{when } 1 \leq j \leq q_{n-2} - s \\ \exp\left(\frac{\pi i}{s} (j - q_{n-2} + s)\right) \cos\left(\frac{\pi}{2s} (j - q_{n-2} + s)\right) & \text{when } q_{n-2} - s < j \leq q_{n-2} \end{cases}$$

$$\mu_j = \begin{cases} 0 & \text{when } 1 \leq j \leq q_{n-2} - s \\ -\exp\left(\frac{\pi i}{s} (j - q_{n-2} + s)\right) \sin\left(\frac{\pi}{2s} (j - q_{n-2} + s)\right) & \text{when } q_{n-2} - s < j \leq q_{n-2}. \end{cases}$$

A careful inspection of the above formulae shows that $W_n^{(k)}$, $1 \leq k \leq a_n$ and \tilde{W}_n are isometries with pairwise orthogonal ranges. This together with (1) implies that W_n is a unitary operator.

For each $n \geq 4$ we shall consider the operators $U_n, V_n \in \mathcal{M}_{q_n}$ defined by:

$$U_n e(n; j) = \begin{cases} e(n; j + 1) & \text{when } 1 \leq j < q_n \\ e(n; 1) & \text{when } j = q_n \end{cases}$$

and

$$V_n e(n; j) = \zeta_n^j e(n; j) \quad \text{where } \zeta_n = \exp\left(2 \pi i \frac{p_n}{q_n}\right).$$

LEMMA 1. For $n \geq 6$ the following inequalities hold:

- (i) $\|W_n^{(k)} U_{n-1} - U_n W_n^{(k)}\| \leq \frac{36 \pi}{q_{n-2}}$
- (ii) $\|W_n^{(k)} V_{n-1} - V_n W_n^{(k)}\| \leq \frac{6 \pi}{q_{n-1}}$

$$(iii) \quad \|\tilde{W}_n U_{n-2} - U_n \tilde{W}_n\| \leq \frac{36\pi}{q_{n-2}}$$

$$(iv) \quad \|\tilde{W}_n V_{n-2} - V_n \tilde{W}_n\| \leq 6\pi \left(\frac{1}{q_{n-1}} + \frac{1}{q_{n-2}} \right).$$

Proof: Applying the operators, to be estimated, to the canonical basis of $C^{q_{n-1}}$ and $C^{q_{n-2}}$ respectively we have:

$$(i') \quad (W_n^{(k)} U_{n-1} - U_n W_n^{(k)}) e(n-1; j) = \begin{cases} (\alpha_{j+1}^{(k)} - \alpha_j^{(k)}) e(n; a(k, j+1)) + (\beta_{j+1}^{(k)} - \beta_j^{(k)}) e(n; b(k, j+1)) + (\gamma_{j+1}^{(k)} - \gamma_j^{(k)}) e(n; c(k, j+1)) & \text{for } 1 \leq j < q_{n-1} \\ \alpha_1^{(k)} e(n; a(k, 1)) + \beta_1^{(k)} e(n; b(k, 1)) + \gamma_1^{(k)} e(n; c(k, 1)) - \alpha_{q_{n-1}}^{(k)} e(n; a(k, q_{n-1}) + 1) - \beta_{q_{n-1}}^{(k)} e(n; b(k, q_{n-1}) + 1) - \gamma_{q_{n-1}}^{(k)} e(n; c(k, q_{n-1}) + 1) & \text{for } j = q_{n-1}. \end{cases}$$

$$(ii') \quad (W_n^{(k)} V_{n-1} - V_n W_n^{(k)}) e(n-1; j) = (\zeta_{n-1}^j - \zeta_n^{a(k,j)}) \alpha_j^{(k)} e(n; a(k, j)) + (\zeta_{n-1}^j - \zeta_n^{b(k,j)}) \cdot \beta_j^{(k)} e(n; b(k, j)) + (\zeta_{n-1}^j - \zeta_n^{c(k,j)}) \gamma_j^{(k)} e(n; c(k, j)).$$

$$(iii') \quad (\tilde{W}_n U_{n-2} - U_n \tilde{W}_n) e(n-2; j) = \begin{cases} (\lambda_{j+1} - \lambda_j) e(n; l(j+1)) + (\mu_{j+1} - \mu_j) e(n; m(j+1)) & \text{for } 1 \leq j < q_{n-2} \\ \lambda_1 e(n; l(1)) + \mu_1 e(n; m(1)) - \lambda_{q_{n-2}} e(n; l(q_{n-2}) + 1) - \mu_{q_{n-2}} e(n; m(q_{n-2}) + 1) & \text{for } j = q_{n-2}. \end{cases}$$

$$(iv') \quad (\tilde{W}_n V_{n-2} - V_n \tilde{W}_n) e(n-2; j) = (\zeta_{n-2}^j - \zeta_n^{l(j)}) \lambda_j e(n; l(j)) + (\zeta_{n-2}^j - \zeta_n^{m(j)}) \mu_j e(n; m(j)).$$

The inequalities:

$$\left| \cos \left(\frac{\pi}{2s} (j+1) \right) - \cos \left(\frac{\pi}{2s} j \right) \right| \leq \frac{\pi}{2s} \leq \frac{2\pi}{q_{n-2}}$$

$$\left| \sin \left(\frac{\pi}{2s} (j+1) \right) - \sin \left(\frac{\pi}{2s} j \right) \right| \leq \frac{\pi}{2s} \leq \frac{2\pi}{q_{n-2}}$$

$$\left| \exp\left(\frac{\pi i}{s}(j+1)\right) \cos\left(\frac{\pi}{2s}(j+1)\right) - \exp\left(\frac{\pi i}{s}j\right) \cos\left(\frac{\pi}{2s}j\right) \right| \leq \frac{3\pi}{2s} \leq \frac{6\pi}{q_{n-2}}$$

$$\left| \exp\left(\frac{\pi i}{s}(j+1)\right) \sin\left(\frac{\pi}{2s}(j+1)\right) - \exp\left(\frac{\pi i}{s}j\right) \sin\left(\frac{\pi}{2s}j\right) \right| \leq \frac{3\pi}{2s} \leq \frac{6\pi}{q_{n-2}}$$

show that $|\alpha_{j+1}^{(k)} - \alpha_j^{(k)}|, |\beta_{j+1}^{(k)} - \beta_j^{(k)}|, |\gamma_{j+1}^{(k)} - \gamma_j^{(k)}|$ ($1 \leq k \leq a_n, 1 \leq j < q_{n-1}$) and $|\lambda_{j+1} - \lambda_j|, |\mu_{j+1} - \mu_j|$ ($1 \leq j < q_{n-2}$) are all bounded by $\frac{6\pi}{q_{n-2}}$.

This implies

$$\|(W_n^{(k)}U_{n-1} - U_nW_n^{(k)})e(n-1; j)\| \leq \frac{18\pi}{q_{n-2}} \quad \text{for } 1 \leq k \leq a_n, 1 \leq j < q_{n-1}$$

and

$$\|(\tilde{W}_nU_{n-2} - U_n\tilde{W}_n)e(n-2; j)\| \leq \frac{12\pi}{q_{n-2}} < \frac{18\pi}{q_{n-2}} \quad \text{for } 1 \leq j < q_{n-2}.$$

To see that both inequalities hold also for $j = q_{n-1}$ and respectively $j = q_{n-2}$, note that $a(k, 1) = c(k, q_{n-1}) + 1$ and $|\alpha_1^{(k)} - \gamma_{q_{n-1}}^{(k)}| < \frac{6\pi}{q_{n-2}}$ for $k > 1$, while for $k = 1, b(1, 1) = c(1, q_{n-1}) + 1$ and $|\lambda_1 - \gamma_{q_{n-1}}^{(1)}| < \frac{6\pi}{q_{n-2}}$, respectively. Also $l(1) = m(q_{n-2}) + 1$ and $|\lambda_1 - \mu_{q_{n-2}}| < \frac{6\pi}{q_{n-2}}$.

This shows that all vectors in (i') and (iii') have norm less than $\frac{18\pi}{q_{n-2}}$. To get the estimates (i) and (iii) remark that the vectors $e(n; a(k, j_1)), e(n; b(k, j_2)), e(n; c(k, j_3)), 1 \leq j_1, j_2, j_3 \leq q_{n-1}$ are pairwise orthogonal for fixed k .

This shows that the operator $W_n^{(k)}U_{n-1} - U_nW_n^{(k)}$ maps the vectors $e(n-1; j)$ ($1 \leq j \leq q_{n-1}$) into pairwise orthogonal vectors. This, now, easily gives

$$\|W_n^{(k)}U_{n-1} - U_nW_n^{(k)}\| \leq \frac{36\pi}{q_{n-2}}.$$

Similarly one gets

$$\|\tilde{W}_nU_{n-2} - U_n\tilde{W}_n\| \leq \frac{36\pi}{q_{n-2}}.$$

Let us now prove the inequalities (ii) and (iv). To this end, note that for every integer m such that $|mq_{n-1} + j| \leq q_n$ we have

$$\begin{aligned} & |\zeta_{n-1}^j - \zeta_n^{mq_{n-1}+j}| = \\ & = \left| \exp\left(2\pi i \frac{p_{n-1}}{q_{n-1}} j\right) - \exp\left(2\pi i \frac{p_n}{q_n} (mq_{n-1} + j)\right) \right| = \\ & = \left| \exp\left(2\pi i \frac{p_{n-1}}{q_{n-1}} (mq_{n-1} + j)\right) - \exp\left(2\pi i \frac{p_n}{q_n} (mq_{n-1} + j)\right) \right| \leq \\ & \leq 2\pi q_n \left| \frac{p_{n-1}}{q_{n-1}} - \frac{p_n}{q_n} \right| = \frac{2\pi}{q_{n-1}} \end{aligned}$$

where the last equality follows from 10.2. Thm. 150 of [8].

Similarly, if m and l are integers such that

$$|mq_{n-1} + lq_{n-2} + j| \leq q_n, \quad |lq_{n-2} + j| \leq q_{n-1},$$

then

$$\begin{aligned} & |\zeta_{n-2}^j - \zeta_n^{mq_{n-1}+lq_{n-2}+j}| \leq \\ & \leq |\zeta_{n-2}^j - \zeta_{n-1}^{lq_{n-2}+j}| + |\zeta_{n-1}^{lq_{n-2}+j} - \zeta_n^{mq_{n-1}+lq_{n-2}+j}| \leq \\ & \leq 2\pi \left(\frac{1}{q_{n-2}} + \frac{1}{q_{n-1}} \right). \end{aligned}$$

These inequalities imply that

$$|\zeta_{n-1}^j - \zeta_n^{a(k,j)}|, |\zeta_{n-1}^j - \zeta_n^{b(k,j)}|, |\zeta_{n-1}^j - \zeta_n^{c(k,j)}|$$

are bounded by $\frac{2\pi}{q_{n-1}}$ for $1 \leq k \leq a_n, 1 \leq j \leq q_{n-1}$, and that

$$|\zeta_{n-2}^j - \zeta_n^{l(j)}|, |\zeta_{n-2}^j - \zeta_n^{m(j)}|$$

are bounded by $2\pi \left(\frac{1}{q_{n-1}} + \frac{1}{q_{n-2}} \right)$ for $1 \leq j \leq q_{n-2}$.

Looking at formulae (ii') and (iv') we have:

$$\| (W_n^{(k)} V_{n-1} - V_n W_n^{(k)}) e(n-1; j) \| \leq \frac{6\pi}{q_{n-1}} \quad \text{for } 1 \leq k \leq a_n, 1 \leq j \leq q_{n-1}$$

$$\| (\tilde{W}_n V_{n-2} - V_n \tilde{W}_n) e(n-2; j) \| \leq 6\pi \left(\frac{1}{q_{n-1}} + \frac{1}{q_{n-2}} \right) \quad \text{for } 1 \leq j \leq q_{n-2}$$

Remarking that the operators in (ii) and (iv) map the canonical basis of $C^{q_{n-1}}$ and respectively $C^{q_{n-2}}$ into pairwise orthogonal vectors, the preceding estimates give (ii) and (iv). Q.E.D.

LEMMA 2.

$$\sum_{n \geq 6} \|\rho_n(U_{n-1} \oplus U_{n-2}) - U_n \oplus U_{n-1}\| < \infty$$

$$\sum_{n \geq 6} \|\rho_n(V_{n-1} \oplus V_{n-2}) - V_n \oplus V_{n-1}\| < \infty.$$

Proof. It is easily seen that the ranges of the operators $W_n^{(k)}U_{n-1} - U_nW_n^{(k)}$ and $W_n^{(j)}U_{n-1} - U_nW_n^{(j)}$ are orthogonal whenever $|k - j| > 6$. Using the preceding lemma this implies that

$$\begin{aligned} & \|W_n(\underbrace{U_{n-1} \oplus \dots \oplus U_{n-1} \oplus U_{n-2}}_{a_n\text{-times}}) - U_nW_n\| \leq \\ & \leq 6 \sup_{1 \leq k \leq a_n} \|W_n^{(k)}U_{n-1} - U_nW_n^{(k)}\| + \|\tilde{W}_nU_{n-2} - U_n\tilde{W}_n\| \leq 7 \cdot \frac{36\pi}{q_{n-2}} \end{aligned}$$

so that

$$\|\rho_n(U_{n-1} \oplus U_{n-2}) - U_n \oplus U_{n-1}\| \leq \frac{300\pi}{q_{n-2}}.$$

The same kind of argument shows that

$$\|\rho_n(V_{n-1} \oplus V_{n-2}) - V_n \oplus V_{n-1}\| \leq \frac{42\pi}{q_{n-1}} + \frac{7\pi}{q_{n-2}}.$$

But $q_n = a_nq_{n-1} + q_{n-2} > 2q_{n-2} \geq \dots \geq 2^{\lfloor \frac{n-1}{2} \rfloor}$ implies the convergence of the series $\sum \frac{1}{q_n}$ which ends the proof. Q.E.D.

The preceding lemma shows that the operators $U_n \oplus U_{n-1} \in A_n \subset A$ and $V_n \oplus V_{n-1} \in A_n \subset A$ are norm-convergent to unitary elements U and respectively V of the AF -algebra A . Since $U_n^*V_nU_n = \exp\left(2\pi i \frac{p_n}{q_n}\right)V_n$ it follows that $U^*VU = \exp(2\pi i\theta)V$. Thus there is a unital $*$ -monomorphism $\rho: A_\theta \rightarrow A$. We thus have proved the following theorem.

THEOREM. *Let $\theta \in (0, 1)$ be an irrational number. There exists a $*$ -monomorphism*

$$\rho: A_\theta \rightarrow A$$

where A is an AF-algebra defined by an inductive limit of finite-dimensional C*-algebras for which the corresponding limit of K-groups is $\dots \rightarrow \mathbf{Z}^2 \xrightarrow{p_n} \mathbf{Z}^2 \rightarrow \dots$ where each \mathbf{Z}^2 is endowed with its natural ordering and

$$\varphi_n = \begin{pmatrix} a_n & 1 \\ 1 & 0 \end{pmatrix}$$

where $[a_1, a_2, \dots, a_n, \dots]$ is the continuous fraction expansion of θ .

By results of E. G. Effros and C. L. Shen [3] we have that $K_0(A)$ is isomorphic to the ordered group $\mathbf{Z} + \theta\mathbf{Z} \subset \mathbf{R}$. In fact, we shall need only the fact that there is a trace-state τ_1 on A such that the range of the induced homomorphism $K_0(A) \rightarrow \mathbf{R}$ is contained in $\mathbf{Z} + \theta\mathbf{Z}$, a fact which can be seen directly as follows. In order to define τ_1 it is sufficient to find positive numbers α_n, β_n corresponding to the values of τ_1 on minimal projections of \mathcal{M}_{a_n} and $\mathcal{M}_{a_{n-1}}$, the two summands of A_n . These numbers must satisfy the relations: $a_n\alpha_n + \beta_n = \alpha_{n-1}$, $\beta_{n-1} = \alpha_n$ and $a_1\alpha_1 + \beta_1 = 1$. Thus it will be sufficient to determine the α_n 's so that $a_1\alpha_1 + \alpha_2 = 1$ and $\alpha_{n+1} + a_n\alpha_n = \alpha_{n-1}$. Taking $\alpha_1 = \theta$ and $\alpha_2 = 1 - a_1\theta$ we can take $\alpha_n = \theta_1 \dots \theta_n$ where $\theta_1 = \theta$ and $\theta_{j+1} = \frac{1}{\theta_j} - a_j$. Since $\alpha_1, \alpha_2 \in \mathbf{Z} + \theta\mathbf{Z}$ we shall have $\alpha_n \in \mathbf{Z} + \theta\mathbf{Z}$ for all $n \in \mathbf{N}$ and thus the range of the homomorphism $K_0(A) \rightarrow \mathbf{R}$ induced by τ_1 is contained in $\mathbf{Z} + \theta\mathbf{Z}$.

Remarking that $\tau_1 \circ \rho = \tau$ we infer that the range of the homomorphism $K_0(A_\theta) \rightarrow \mathbf{R}$ induced by τ is contained in $\mathbf{Z} + \theta\mathbf{Z}$. Thus, using Rieffel's result mentioned in the introduction we have:

COROLLARY 1. *Let $\psi: K_0(A_\theta) \rightarrow \mathbf{R}$ be the homomorphism induced by τ . Then $\psi(K_0(A_\theta)) = \mathbf{Z} + \theta\mathbf{Z}$.*

Let now $\theta_1, \theta_2 \in (0, 1)$ be irrational numbers and remark that $\mathbf{Z} + \theta_1\mathbf{Z} = \mathbf{Z} + \theta_2\mathbf{Z}$ as subsets of \mathbf{R} if and only if $\theta_1 \in \{\theta_2, 1 - \theta_2\}$. This together with the isomorphism $A_{\theta_2} \simeq A_{1-\theta_2}$ gives:

COROLLARY 2. *Let $\theta_1, \theta_2 \in (0, 1)$ be irrational numbers. Then A_{θ_1} and A_{θ_2} are isomorphic if and only if $\theta_1 \in \{\theta_2, 1 - \theta_2\}$.*

Added in Proof

1) After this paper had been circulated as a preprint (INCREST-preprint no. 45/1979) several advances have been made concerning K-groups of crossed products (see the Added in Proof of our paper "Exact sequences for K-groups and Ext-groups of certain cross-product C*-algebras", *J. Operator Theory* 4 (1980), 93-118).

2) The authors thank Professor A. M. Vershik for pointing out that the imbedding result of the present paper (used together with a non-commutative Weyl-von Neumann type theorem obtained by the second author) provides an affirmative answer to a problem raised by Professor

Vershik concerning approximability in norm of the pair of unitaries corresponding to the usual representation in $L^2(\mathbb{T})$ of A_θ , by pairs of unitaries contained in finite-dimensional C^* -algebras (see Zapiski nauchnih seminarov LOMI vol. 81, "99 unsolved problems of linear and complex analysis", pp. 77- 81).

3) We would like to mention that our construction for the imbedding $A_\theta \hookrightarrow A$ is related to I. D. Berg's technique for the approximation of weighted shifts.

REFERENCES

1. BRATTELI, O., Inductive limits of finite-dimensional C^* -algebras, *Trans. Amer. Math. Soc.*, **171** (1972), 195--234.
2. CONNES, A., Sur la théorie non-commutative de l'intégration, Preprint.
3. EFFROS, E. G.; SHEN, C. L., Approximately finite C^* -algebras and continued fractions, Preprint.
4. EFFROS, E. G.; SHEN, C. L., Dimension groups and finite difference equations, *J. Operator Theory*, **2** (1979), 215--231.
5. EFFROS, E. G.; ; HANDELMAN, D.; SHEN, C. L., Dimension groups and their affine representations, Preprint.
6. EFFROS, P. G.; SHEN, C. L., The geometry of finite rank dimension group, Preprint.
7. ELLIOTT, G., On the classification of inductive limits of sequences of semi-simple finite dimensional algebras, *J. Algebra*, **38** (1976), 29--44.
8. HARDY, G. H.; WRIGHT, E. M., *An introduction to the theory of numbers*, 1954, Oxford.
9. RIEFFEL, M. A., *Irrational rotation C^* -algebras*, short communication at the International Congress of Mathematicians, 1978.
10. SHEN, C. L., A note on the automorphism group of simple dimension groups, Preprint.
11. SHEN, C. L., On the classification of the ordered groups associated with the approximately finite dimensional C^* -algebras, Preprint.

M. PIMSNER and D. VOICULESCU
Department of Mathematics,
National Institute for Scientific and
Technical Creation,
Bd. Păcii 220, 79622 Bucharest,
Romania

Received November 26, 1979.