

## ONE-PARAMETER AUTOMORPHISM GROUPS OF THE HYPERFINITE TYPE $II_1$ FACTOR

YASUYUKI KAWAHIGASHI

### 0. INTRODUCTION

In this paper we show the uniqueness, up to cocycle conjugacy, of a one-parameter automorphism group of the hyperfinite type  $II_1$  factor  $\mathcal{R}$ , which fixes a Cartan subalgebra of  $\mathcal{R}$  elementwise and has the Connes spectrum  $\hat{\mathbf{R}}$ . This result is valid for any separable locally compact abelian group  $G$  instead of  $\mathbf{R}$ . As its application, we also show the uniqueness, up to cocycle conjugacy, of an almost periodic prime action  $\alpha$  of a separable locally compact abelian group on the hyperfinite type  $II_1$  factor  $\mathcal{R}$  with  $(\mathcal{R}^\alpha)' \cap \mathcal{R} = \mathbf{C}I$ , and the uniqueness, up to cocycle conjugacy, of a quasi-free one-parameter automorphism group of  $\mathcal{R}$  arising from the CAR  $C^*$ -algebra, which has the Connes spectrum  $\mathbf{R}$ . The ways of computation of the Connes spectrum in terms of the asymptotic range are also given. Several examples are shown to be identical up to cocycle or stably conjugacy as an application.

In this classification problem of group actions on the hyperfinite type  $II_1$  factor, non-compact continuous group have not been studied well. Thus we study the classification problem for the real number group  $\mathbf{R}$ . Note that the uniqueness of the injective type  $III_1$  factor, which was finally solved by [8], is equivalent to the uniqueness, up to conjugacy, of the trace scaling one-parameter automorphism group,  $\text{tr} \circ \alpha_t = e^{-t} \text{tr}$ , of the hyperfinite type  $II_\infty$  factor  $\mathcal{R}_{0,1}$  by [23]. The trace preserving cases are still open.

We solved the classification problem for an action  $\alpha$  of  $\mathbf{R}$  up to stable conjugacy in the previous paper [13] for the cases  $\Gamma(\alpha) \neq \mathbf{R}$ . In Section 1, we will deal with the case  $\Gamma(\alpha) = \mathbf{R}$  under the condition that the action  $\alpha$  fixes a Cartan subalgebra of  $\mathcal{R}$ . If an action fixes a Cartan subalgebra elementwise, we can write down the explicit form of this type of action by the works of [7] and [5], and we will classify this type

of actions. We will use the technique of  $T$ -array in [14], and reduce the general cases to infinite tensor product type actions. The results in this section are stated for a general separable locally compact abelian group  $G$ . The method in this section can also be applied to the hyperfinite type  $\text{II}_\infty$  factor.

In Section 2 we show uniqueness, up to cocycle conjugacy, of an almost periodic prime action  $\alpha$  of a separable locally compact abelian group on the hyperfinite type  $\text{II}_1$  factor  $\mathcal{R}$  with  $(\mathcal{R}^\alpha)' \cap \mathcal{R} = \mathbf{C}I$  as an application of the result in Section 1. This type of actions were studied by [25].

In Section 3 we will use the construction in Section 1 and its modification to show that all the ergodic flows actually occur as  $\hat{\alpha}$  on  $\mathcal{Z}(\mathcal{M} \rtimes_\alpha \mathbf{R})$ , which was used as complete invariants together with the type of the crossed product algebra for the classification of an action  $\alpha$  of  $\mathbf{R}$  on the hyperfinite type  $\text{II}_1$  or  $\text{II}_\infty$  factor  $\mathcal{M}$  with  $\Gamma(\alpha) \neq \mathbf{R}$  in [13]. We also show a one-parameter automorphism group  $\alpha$  has the trivial relative commutant property  $\mathcal{R}' \cap \mathcal{R} \rtimes_\alpha \mathbf{R} = \mathbf{C}I$  if  $\alpha$  fixes a Cartan subalgebra of  $\mathcal{R}$  and  $\Gamma(\alpha) = \mathbf{R}$ . We show examples of a one-parameter automorphism with  $\Gamma(\alpha) = \mathbf{R}$  at the end of this section.

In Section 4 we use the result in Section 1 for quasi-free actions of  $\mathbf{R}$  on  $\mathcal{R}$ , which are weak extensions of a one-parameter automorphism group on the CAR  $C^*$ -algebra coming from the Bogoliubov automorphism given by a one-parameter unitary group on a separable Hilbert space. We reduce these actions to the above type of actions by expansionals in [1]. As an example, we can apply this result to “CAR-flow” which is a type  $\text{II}_1$  factor automorphism group version of the endomorphism semigroup of  $\mathcal{L}(\mathcal{H})$  in [21].

We would like to thank Prof. K. Schmidt for calling our attention to [17], Prof. R. Powers for raising our interest in the subject of Section 4, and Prof. M. Takesaki for numerous suggestions and successive encouragement.

## 1. UNIQUENESS RESULT FOR LOCALLY COMPACT ABELIAN GROUPS

We study actions of a locally compact abelian group  $G$  on the hyperfinite type  $\text{II}_1$  factor  $\mathcal{R}$  defined as follows. (Though our main interest lies in the case  $G = \mathbf{R}$ , the results in this section are valid for more general cases.) Let  $T$  be an ergodic measure preserving transformation on a measure space  $(X, \mu)$ ,  $\mu(X) = 1$ . Then  $\sigma \in \text{Aut}(L^\infty(X, \mu))$  is defined by  $\sigma(\varphi)(x) = \varphi(T^{-1}x)$  for  $\varphi \in L^\infty(X, \mu)$ . The group measure space construction  $L^\infty(X, \mu) \rtimes_\sigma \mathbf{Z}$  gives us  $\mathcal{R}$ . Let  $u$  be the implementing unitary for this crossed product algebra. We denote  $L^\infty(X, \mu)$  by  $\mathcal{A}$  in the following. For a separable locally compact abelian group  $G$ , take a measurable function  $h$  from

$X$  to  $\hat{G}$ , and we define an action  $\alpha_t$  of  $G, t \in G$  by

$$\begin{cases} \alpha_t(\varphi) = \varphi, & \text{for } \varphi \in L^\infty(X, \mu) \\ \alpha_t(u) = \langle t, h(x) \rangle u, \end{cases}$$

where  $\langle t, h(x) \rangle$  denotes the duality pairing of  $t \in G$  and  $h(x) \in \hat{G}$  for  $x \in X$ . Note that if  $\alpha$  is an action of  $G$  which fixes a Cartan subalgebra  $\mathcal{R}$  elementwise, then  $\alpha$  is of this form. (See Definition 3.1, Theorem 1, Theorem 5 in [7] and Theorem 10 in [5].) In this section,  $\alpha$  will denote this action. Every set in the following is assumed to be measurable.

We use a groupoid  $X \rtimes_T \mathbb{Z}$  for which the multiplication is defined by  $(x, n) \cdot (T^{-n}x, m) = (x, n + m)$ , where  $x \in X, n, m \in \mathbb{Z}$ .

**DEFINITION 1.1.** For the above measurable function  $h$  and an ergodic transformation  $T$  on  $X$ , we denote by  $r(h, T)$  the asymptotic range  $r^*(c)$  (Definition 8.2 in [6]) for the following cocycle on the groupoid  $X \rtimes_T \mathbb{Z}$ .

$$c(x, n) = \begin{cases} h(x) + h(T^{-1}x) + \dots + h(T^{-n+1}x), & \text{if } n > 0, x \in X, \\ 0, & \text{if } n = 0, x \in X, \\ h(Tx) + h(T^2x) + \dots + h(T^{-n}x), & \text{if } n < 0, x \in X. \end{cases}$$

The asymptotic range  $r^*(c)$  is a closed subgroup of  $\hat{G}$  by Proposition 8.5 in [6].

For the Connes spectrum  $\Gamma(\alpha)$  (see Définition 2.2.1 in [3]), we get the following, as expected. (See Proposition 2.11 in [7].)

**PROPOSITION 1.2.** For the above action  $\alpha, \Gamma(\alpha) = r(h, T)$ .

*Proof.* By IV.5.4. in [24] and  $\mathcal{Z}(\mathcal{R}^\infty) \subseteq \mathcal{A} \subseteq \mathcal{R}^\infty$  we get

$$\begin{aligned} \Gamma(\alpha) &= \bigcap \{ \text{Sp}(\alpha^e) \mid e \in \text{Proj}(\mathcal{R}^\infty) \} \subseteq \bigcap \{ \text{Sp}(\alpha^e) \mid e \in \text{Proj}(\mathcal{A}) \} \subseteq \\ &\subseteq \bigcap \{ \text{Sp}(\alpha^e) \mid e \in \text{Proj}(\mathcal{Z}(\mathcal{R}^\alpha)) \} = \Gamma(\alpha). \end{aligned}$$

(Here the symbol  $\mathcal{Z}$  means the center.) Thus  $\lambda \in \Gamma(\alpha)$  if and only if  $\lambda \in \text{Sp}(\alpha^e)$  for every  $e = \chi_B \in \mathcal{A}$ , where  $B \subseteq X, \mu(B) > 0$ .

Suppose  $\lambda \in r(h, T)$ . Choose an arbitrary  $B \subseteq X, \mu(B) > 0$ , and set  $e = \chi_B \in \mathcal{A}$ . We have to show  $\hat{f}(\lambda) = 0$  if we have an  $f \in L^1(G)$  such that

$$(*) \quad \int_G f(-t) \alpha_t^e(y) dt = 0, \quad \text{for every } y \in \mathcal{R}_e.$$

Assume  $\hat{f}(\lambda) \neq 0$ , and take an open neighborhood  $U \subseteq \hat{G}$  of  $\lambda$  so that  $\hat{f} \neq 0$  on  $U$ . Now by the definition of the asymptotic range, there exists an integer  $n$  such that we have  $\mu(B') > 0$  for

$$B' = \{ x \in B \mid T^{-n}x \in B, c(x, n) \in U \},$$

where we used the cocycle  $c$  as in Definition 1.1. Take  $y = eu^n e$  in (\*). We have

$$\int_G f(-t) \chi_B \alpha_t(u^n) \chi_B dt = \int_G f(t) \chi_B(x) \overline{(t, c(x, n))} \chi_{T^{-n}B}(x) dt \cdot u^n = 0.$$

This implies

$$\int_G f(t) \overline{(t, c(x, n))} dt = 0, \quad \text{for almost all } x \in B'.$$

This contradicts  $c(x, n) \in U$  for  $x \in B'$  and  $\hat{f} \neq 0$  on  $U$ .

Conversely, assume  $\lambda \in \text{Sp}(\alpha^e)$  for every  $e = \chi_B$ ,  $B \subseteq X$ ,  $\mu(B) > 0$ . Suppose  $\lambda \notin r(h, T)$ . Then there exist  $B \subseteq X$ ,  $\mu(B) > 0$ , and a neighborhood  $U \subseteq \hat{G}$  of  $\lambda$  such that

$$(**) \quad \mu(\{x \in B \mid T^{-n}x \in B, c(x, n) \in U\}) = 0, \quad \text{for every integer } n.$$

Take an  $f \in L^1(G)$  such that  $\text{supp}(\hat{f}) \subseteq U$  and  $\hat{f}(\lambda) \neq 0$ . Then for every integer  $n$  and  $\varphi \in L^\infty(X, \mu)$ , we have

$$\int_G f(-t) \alpha_t(e\varphi u^n e) dt = \int_G f(t) \chi_B \varphi(t, c(x, n)) \chi_{T^{-n}B} dt \cdot u^n.$$

But the right hand side of this is 0 because of (\*\*) and  $\text{supp}(\hat{f}) \subseteq U$ . Thus by the definition of  $\text{Sp}(\alpha^e)$ , we get  $\hat{f}(\lambda) = 0$ , which contradicts the construction of  $f$ .

Q.E.D.

We get following for the Poincaré flow. (See Definition 8.1 in [6].)

**PROPOSITION 1.3.** *For the above action  $\alpha$ , the flow on  $Z(\mathcal{R} \rtimes_\alpha G)$  given by the dual action  $\hat{\alpha}$  of  $\hat{G}$  is the Poincaré flow of the cocycle  $c$ .*

*Proof.* By considering the dual action  $\hat{\alpha}$  on  $Z(L^\infty(X, \mu) \rtimes_\sigma \mathbb{Z} \rtimes_\alpha G)$ , it follows easily from the definition of the Poincaré flow. Q.E.D.

We are interested in the case  $r(h, T) = \Gamma(\alpha) = \hat{G}$ . Thus in the rest of this section, we assume this equality. In the following we use the notation  $[T]$  for the full group of  $T$ .

**DEFINITION 1.4.** For  $S \in [T]$  and a measurable function  $h$  from  $X$  to  $\hat{G}$ , we define

$$F(h; x, S) = \begin{cases} h(x) + h(T^{-1}x) + \cdots + h(T^{-n+1}x), & \text{if } n > 0, \\ 0, & \text{if } n = 0 \\ h(Tx) + h(T^2x) + \cdots + h(T^{-n}x), & \text{if } n < 0, \end{cases}$$

on the set  $\{x \in X \mid Sx = T^{-n}x\}$ .

LEMMA 1.5. For every  $A, B \subseteq X$ ,  $\mu(A) = \mu(B) > 0$ ,  $\lambda \in \hat{G}$  and a neighborhood  $W \subseteq \hat{G}$  of  $\lambda$ , there exists  $S \in [T]$  such that  $S(A) = B$ ,  $F(h; x, S) \in W$  for almost all  $x \in A$ .

*Proof.* Because  $T$  is ergodic, there exist  $A_1 \subseteq A$ ,  $B_1 \subseteq B$ ,  $\mu(A_1) = \mu(B_1) > 0$  and an integer  $n$  such that  $T^{-n}A_1 = B_1$ . By considering  $c(x, n)$  on  $A_1$ , there exists  $A_2 \subseteq A_1$ ,  $B_2 \subseteq B_1$ ,  $\mu(A_2) = \mu(B_2) > 0$ ,  $\lambda' \in \hat{G}$ , an open neighborhood  $W' \subseteq \hat{G}$  of  $\lambda'$ , and an open neighborhood  $W'' \subseteq \hat{G}$  of  $\lambda - \lambda'$  such that  $c(x, n) \in W'$  for almost all  $x \in A_2$  and  $W' + W'' \subseteq W$  because  $\hat{G}$  is second countable. Then by the definition of the asymptotic range, there exist  $B_3, B_4 \subseteq B_2$ ,  $\mu(B_3) = \mu(B_4) > 0$ , and an integer  $m$  such that  $T^{-m}B_3 = B_4$  and  $c(x, m) \in W''$  for almost all  $x \in B_3$ . Now set  $A_3 = T^n B_3$ . Then we have  $c(x, n + m) \in W$  for almost all  $x \in A_3$ .

Now let  $\mathcal{F}$  be the set of families  $\{A_i, B_i\}_{i \in I}$ , where  $\mu(A_i) = \mu(B_i) > 0$ ,  $A_i$ 's are mutually disjoint subsets of  $A$ ,  $B_i$ 's are mutually disjoint subsets of  $B$ , and for each  $i \in I$  there exists an integer  $n_i$  such that  $T^{-n_i}A_i = B_i$  and  $c(x, n_i) \in W$  for almost all  $x \in A_i$ . Consider the usual order on  $\mathcal{F}$ , then it is inductively ordered. Thus take a maximal  $\{A_i, B_i\}_{i \in I}$  in  $\mathcal{F}$ , then  $\mu(\cup A_i) = \mu(\cup B_i)$ . If  $\mu(A - \cup A_i) = \mu(B - \cup B_i) > 0$ , then we can find another  $A'$  and  $B'$  by applying the above argument to  $A - \cup A_i$ ,  $B - \cup B_i$ , which contradicts the maximality of  $\{A_i, B_i\}_{i \in I}$ . Now  $A = \cup A_i$ , and  $B = \cup B_i$ , thus we are done. (The transformation  $S$  is defined to be  $T^{-n_i}$  on  $A_i$ .)

Q.E.D.

While this Lemma 1.5 is similar to Lemma 2.7 in [14], the important difference is that  $\lambda$  is arbitrary here.

Take an action  $\beta$  of  $G$  on  $\mathcal{R}$  of the form  $\beta_t = \bigotimes_{j=1}^{\infty} \text{diag}(\langle t, \nu_0^j \rangle, \dots, \langle t, \nu_{N_j}^j \rangle)$ , where  $\text{diag}(\langle t, \nu_0^j \rangle, \dots, \langle t, \nu_{N_j}^j \rangle)$  stands for the  $(N_j + 1) \times (N_j + 1)$  diagonal matrix with diagonal entries  $\langle t, \nu_0^j \rangle, \dots, \langle t, \nu_{N_j}^j \rangle$ , and  $\nu_i^j$ 's are in  $\hat{G}$ . We say this action is of the infinite tensor product type. In this expression, we may assume  $\nu_0^j = 0$  for all  $j$ , hence we assume this in this section, and fix  $\beta$ . We will prove the following theorem on the uniqueness of actions up to cocycle conjugacy. (See p.215 of [10] for the definition of cocycle conjugacy.)

THEOREM 1.6. If an action  $\alpha$  of a locally compact separable abelian group  $G$  on the hyperfinite type II<sub>1</sub> factor  $\mathcal{R}$  fixes a Cartan subalgebra elementwise and  $\Gamma(\alpha) = \hat{G}$ , and another action  $\beta$  is of the infinite tensor product type, then  $\alpha$  is cocycle conjugate to an action of the infinite tensor product type, and  $\alpha \otimes \beta$  is cocycle conjugate to  $\alpha$ .

Note that the infinite tensor product type actions are particular cases of the actions in this theorem.

We need some lemmas for the proof of this theorem. We will use the technique

of  $T$ -array of Krieger. (See p.166 in [14] and V.5 in [24] for definitions and notations.) In our convention here, we assume  $U(a, b)Z(a) = Z(b)$ , the index set  $A$  is finite, and  $\bigcup_{a \in A} Z(a) = X$  for a  $T$ -array  $\mathcal{A} = \{Z(a), U(a, b) \mid a, b \in A\}$ . We use the notation  $\partial k(x) = k(x) - k(T^{-1}x)$  for a measurable function  $k$  from  $X$  to  $\hat{G}$ .

LEMMA 1.7. Suppose a  $T$ -array  $\mathcal{A}_1 = \{Z_1(a), U_1(a, b) \mid a, b \in A_1\}$ ,  $B_1, \dots, B_m \subseteq X$ , a measurable function  $h_1$  from  $X$  to  $\hat{G}$ ,  $\varepsilon > 0$ , and an open neighborhood  $W \subseteq \hat{G}$  of 0 are given. Moreover, we assume that  $F(h_1; x, U_1(a, b))$  is an almost everywhere constant function on  $Z_1(b)$ . Then there exists an integer  $n_0$  such that for every integer  $n \geq n_0$ , and  $\lambda_0, \dots, \lambda_{n-1} \in \hat{G}$ , where  $\lambda_0 = 0$ , there exist an extension  $T$ -array of  $\mathcal{A}_1$

$$\mathcal{A}_2 = \{Z_2(a), U_2(a, b) \mid a, b \in A_2 = A_1 \times \mathbb{Z}_n\}$$

and a measurable function  $h_2$  from  $X$  to  $\hat{G}$  such that

$$h_2(x) \in W, \quad \text{for almost all } x \in X,$$

$$B_k \overset{\varepsilon}{\in} \mathcal{B}\{Z_2(a) \mid a \in A_2\}, \quad \text{for every } 1 \leq k \leq m,$$

$$F(h_1; x, U_1(a, b)) = F(h_1 + \partial H_2; x, U_1(a, b)), \quad \text{for almost all } x \in X,$$

$$F(h_1 + \partial h_2; x, U_2((a, j), (a, 0))) = \lambda_j, \quad a \in A_1, \quad \text{for almost all } x \in Z_2(a, 0).$$

*Proof.* Take  $n_0$  as in Lemma V.5.7 in [24], and for  $n \geq n_0$ , take  $Z_2(a, j)$  for  $(a, j) \in A_1 \times \mathbb{Z}_n$  as in the proof of Lemma V.5.7 in [24] so that  $B_k \overset{\varepsilon}{\in} \mathcal{B}\{Z_2(a) \mid a \in A_2\}$  where the right hand side means the  $\sigma$ -algebra generated by  $Z_2(a)$ ,  $a \in A_2$ . Fix  $a_0 \in A$ . Now take  $U_2((a_0, j), (a_0, 0)) \in [T]$  by Lemma 1.5 so that

$$F(h_1; x, U_2((a_0, j), (a_0, 0))) \in \lambda_j + W, \quad \text{for almost all } x \in Z_2(a_0, 0).$$

(For  $j = 0$ , take  $U_2((a_0, 0), (a_0, 0)) = \text{id}$ .) Now define

$$U_2((a_0, i), (a_0, j)) = U_2((a_0, i), (a_0, 0))U_2((a_0, j), (a_0, 0))^{-1},$$

and extend this as usual. (See V.5.6 in [24].) We define

$$h_2(x) = F(h_1; U_2((a, 0), (a, j))x, U_2((a, j), (a, 0))) - \lambda_j$$

on  $Z_2(a, j)$ ,  $(a, j) \in A_1 \times \mathbb{Z}_n$ . Then  $h_2(x) \in W$  for almost all  $x \in X$ . For almost all  $x \in Z(a, 0)$ , we have

$$F(h_1 + \partial h_2; x, U_2((a, j), (a, 0))) =$$

$$\begin{aligned}
&= h_1(x) + h_1(T^{-1}x) + \cdots + h_1(T^{-n+1}x) + h_2(x) - h_2(T^{-n}x) = \\
&= h_1(x) + h_1(T^{-1}x) + \cdots + h_1(T^{-n+1}x) - F(h_1; x, U_2((a, j), (a, 0))) + \lambda_j = \lambda_j,
\end{aligned}$$

where  $n$  is given by  $U_2((a, j), (a, 0))x = T^{-n}x$ . Thus by construction of  $h_2$ , we have the desired equalities.

We use the notation  $\text{Orb}_U(x) = \{U(a, b)x \mid a, b \in A\}$  for a  $T$ -array  $\mathcal{A} = \{Z(a), U(a, b) \mid a, b \in A\}$  and  $x \in X$ .

LEMMA 1.8. For a given  $T$ -array  $\mathcal{A}_1 = \{Z_1(a), U_1(a, b) \mid a, b \in A_1\}$ ,  $\varepsilon > 0$ , an open neighborhood  $W \subseteq \hat{G}$  of 0, and a measurable function  $h_1$  from  $X$  to  $\hat{G}$  such that  $F(h_1; U_1(a, b), x)$  is an almost everywhere constant function on  $Z_1(b)$ , there exist an integer  $n$ , an extension  $T$ -array  $\mathcal{A}_2 = \{Z_2(a), U_2(a, b) \mid a, b \in A_2 = A_1 \times \mathbb{Z}_n\}$ , a measurable function  $h_2$  from  $X$  to  $\hat{G}$ , and  $\rho_0, \dots, \rho_{n-1} \in \hat{G}$  such that

$$\mu(\{x \in X \mid Tx \notin \text{Orb}_{U_2}(x)\}) < \varepsilon,$$

$$\mu(\{x \in X \mid h_2(x) \in W\}) > 1 - \varepsilon,$$

$$F(h_1; x, U_1(a, b)) = F(h_1 + \partial h_2; x, U_1(a, b)), \quad \text{for almost all } x \in Z_1(b),$$

$$F(h_1 + \partial h_2; x, U_2(a, j), (a, 0)) = \rho_j, \quad a \in A_1, \quad \text{for almost all } x \in Z_2(a, 0).$$

*Proof.* First, take an extension  $T$ -array  $\mathcal{A}'_1 = \{Z'_1(a), U'_1(a, b) \mid a, b \in A'_1 = A_1 \times \mathbb{Z}_m\}$  for some integer  $m$  by Lemma V.5.8 in [24] such that

$$\mu(\{x \in X \mid Tx \notin \text{Orb}_{U'_1}(x)\}) < \varepsilon.$$

We make an extension of this by technique on p.168 in [14]. Fix  $a_0 \in A_1$ . Then there exist  $E \subseteq Z'_1(a_0, 0)$ , an open neighborhood  $W' \subseteq \hat{G}$  of 0, and  $\rho'_0, \dots, \rho'_{m-1} \in \hat{G}$  such that  $\mu(E) > 0$ ,  $W' + W' \subseteq W$ , and

$$F(h_1; x, U'_1(a_0, j), (a_0, 0)) \in \rho'_j + W', \quad 0 \leq j \leq m-1, \quad \text{for almost all } x \in E.$$

By maximality argument, we can find a family of mutually disjoint sets  $\{E'_i\}_{i \in \mathbb{N}}$  in  $Z'_1(a_0, 0)$  and elements  $\{\rho'_{i,j}\}_{i \in \mathbb{N}, 0 \leq j \leq m-1}$  of  $\hat{G}$  such that  $\rho'_{i,0} = 0$ ,  $\mu(E'_i) > 0$ ,  $\mu(Z'_1(a_0, 0) - \bigcup_{i \in \mathbb{N}} E'_i) = 0$  and

$$F(h_1; x, U'_1((a_0, j), (a_0, 0))) \in \rho'_{i,j} + W', \quad \text{for all } j \text{ and almost all } x \in E'_i.$$

Take  $l_0 \in \mathbb{N}$  such that  $\mu(\bigcup_{i \geq l_0} E'_i) < \varepsilon \mu(Z'_1(a_0, 0))/2$ . By approximating  $\bigcup_{i \geq l_0} E'_i$  and  $E'_i$ 's ( $0 \leq i \leq l_0 - 1$ ) by unions of smaller sets, we get integers  $l_1, l$ , a family  $\{E_i\}_{0 \leq i \leq l-1}$

of mutually disjoint sets in  $Z'_1(a_0, 0)$  and elements  $\{\rho_{i,j}\}_{0 \leq i \leq l-1, 0 \leq j \leq m-1}$  of  $\hat{G}$  such that

$$\begin{aligned} \rho_{i,0} &= 0, \\ \mu(E_i) &= \mu(E'_i), \\ \mu \left( Z'_1(a_0, 0) - \bigcup_{0 \leq i \leq l-1} E_i \right) &= 0, \\ \mu \left( \bigcup_{l_1 \leq i \leq l-1} E_i \right) &\leq \varepsilon \mu(Z'_1(a_0, 0)), \end{aligned}$$

$$F(h_1; x, U'_1((a_0, j), (a_0, 0))) \in \rho_{i,j} + W', \quad 0 \leq i \leq l_1 - 1 \quad \text{for almost all } x \in E_i.$$

We define  $Z_2(a_0, j, i) = U'_1((a_0, j), (a_0, 0))E_i$ , choose  $U_2((a_0, 0, i), (a_0, 0, 0)) \in [T]$  such that

$$\begin{aligned} U_2((a_0, 0, i), (a_0, 0, 0))Z_2(a_0, 0, 0) &= Z_2(a_0, 0, i), \\ U_2((a_0, 0, 0), (a_0, 0, 0)) &= \text{id}, \end{aligned}$$

$$F(h_1; x, U_2((a_0, 0, i), (a_0, 0, 0))) \in W', \quad \text{for almost all } x \in Z_2(a_0, 0, 0),$$

by Lemma 1.5. Now we can define

$$U_2((a_0, j, i), (a_0, 0, 0)) = U'_1((a_0, j), (a_0, 0))U_2((a_0, 0, i), (a_0, 0, 0)),$$

and extend this as usual. Now we define

$$h_2(x) = F(h_1; U_2((a_0, 0, 0), (a_0, j, i))x, U_2((a_0, j, i), (a_0, 0, 0))) - \rho_{i,j}$$

on  $Z_2(a_0, j, i)$ . Thus this  $h_2(x)$  is defined on  $Z_1(a_0)$ . We extend this to the entire set  $X$  by  $h_2(x) = h_2(U_1(a_0, a)x)$  on  $Z_1(a)$ . Then we know that  $\mu(\{x \in X \mid h_2(x) \in W\}) > 1 - \varepsilon$ . Set  $n = lm$ . Because two equalities for  $F$  are proved as in the proof of Lemma 1.7,  $\mathcal{A}_2 = \{Z_2(a), U_2(a, b) \mid a, b \in \mathcal{A}_2 = \mathcal{A}_1 \times \mathbb{Z}_n\}$ ,  $\rho$ 's and  $h_2$  satisfy the desired properties. Q.E.D.

Now we can prove Theorem 1.6.

*Proof of Theorem 1.6.* Let  $\tilde{\beta}$  be the infinite tensor product of copies of  $\beta$ . Because  $\tilde{\beta}$  is also of the infinite tensor product type, we represent this as

$$\tilde{\beta} = \bigoplus_{j=0}^{\infty} \text{Ad}(\text{diag}(\langle t, \lambda_0^j \rangle, \dots, \langle t, \lambda_{l_j}^j \rangle)), \quad \lambda_i^j \in \hat{G}.$$



Let  $\{B_n\}_{n \in \mathbf{N}}$  be a sequence of Borel sets which generates the  $\sigma$ -algebra of  $X$ . Choose a sequence  $\{W_j\}_{j \in \mathbf{N}}$  of open neighborhoods of 0 in  $\hat{G}$  such that  $\sum_{j=0}^{\infty} \lambda_j$  always converges for an arbitrary sequence  $\{\lambda_j\}_{j \in \mathbf{N}}$ ,  $\lambda_j \in W_j$ . We can construct a sequence of  $T$ -arrays,  $\mathcal{A}_0^1, \mathcal{A}_0^2, \mathcal{A}_1^1, \mathcal{A}_1^2, \dots$ , a sequence of measurable functions from  $X$  to  $\hat{G}$ ,  $\{h_j^1, h_j^2\}_{j \in \mathbf{N}}$ , a sequence of integers  $\{n_j^1, n_j^2\}_{j \in \mathbf{N}}$ , a sequence of elements  $\{\rho_{j,k}^1, \rho_{j,k}^2\}_{j \in \mathbf{N}, 0 \leq k \leq n_j-1}$  of  $\hat{G}$ , ( $\rho_{j,0}^1 = \rho_{j,0}^2 = 0$ ), and a strictly increasing sequence of integers  $\{m_j\}_{j \in \mathbf{N}}$ , ( $m_0 = 0$ ), by applying Lemma 1.7 and Lemma 1.8 alternately so that the following conditions are satisfied:

$$(1) \quad \mathcal{A}_j^i = \{Z_j^i(a), U_j^i(a, b) \mid a, b \in A_j^i\}, \quad i = 1, 2, j \in \mathbf{N},$$

$$(2) \quad \mathcal{A}_j^2 \text{ is an extension of } \mathcal{A}_j^1,$$

$$(3) \quad \mathcal{A}_{j+1}^1 \text{ is an extension of } \mathcal{A}_j^2,$$

$$(4) \quad \mathcal{A}_j^1 = \mathbf{Z}_{n_1^1} \times \mathbf{Z}_{n_2^1} \times \cdots \times \mathbf{Z}_{n_j^1},$$

$$(5) \quad \mathcal{A}_j^2 = \mathcal{A}_j^1 \times \mathbf{Z}_{n_k^2},$$

$$(6) \quad B_k \stackrel{1/2^j}{\in} \mathcal{B}\{Z_j^1(a) \mid a \in A_j^1\}, \quad k \leq j,$$

$$(7) \quad \mu(\{x \in X \mid Tx \in \text{Orb}_{U_j^2}(x)\}) > 1 - 1/2^j,$$

$$(8) \quad h_j^1(x) \in W_j, \quad \text{for almost all } x \in X,$$

$$(9) \quad \mu(\{x \in X \mid h_j^2(x) \in W_j\}) \geq 1 - 1/2^j,$$

$$(10) \quad F(h + \partial(h_0^1 + h_0^2 + h_1^1 + h_1^2 + \cdots + h_j^1); x, U_j^1((a, k), (a, 0))) = \rho_{j,k}^1,$$

for  $a \in A_{j-1}^2, 0 \leq k \leq n_j^1 - 1$ , and almost all  $x \in Z_k^1(a, 0)$ ,

$$(11) \quad F(h + \partial(h_0^1 + h_0^2 + h_1^1 + h_1^2 + \cdots + h_j^2); x, U_j^2((a, k), (a, 0))) = \rho_{j,k}^2,$$

for  $a \in A_j^1$ ,  $0 \leq k \leq n_j^2 - 1$ , and almost all  $x \in Z_k^2(a, 0)$ ,

$$(12) \quad \text{diag}(\langle t, \rho_{j,0}^1 \rangle, \dots, \langle t, \rho_{j,n_j^1}^1 \rangle) = \bigotimes_{n=m_j}^{m_{j+1}-1} \text{diag}(\langle t, \lambda_0^n \rangle, \dots, \langle t, \lambda_{n_j}^n \rangle).$$

Note that there exists a measurable function  $h' = \sum_{j=0}^{\infty} (h_j^1 + h_j^2)$  on  $X$  by (8) and

(9). By (2), (3), (4), (5), (6) and (7),  $L^\infty(X, \mu) \rtimes_\sigma \mathbb{Z}$  is isomorphic to  $L^\infty(\prod_{j=0}^{\infty} (\mathbb{Z}_{n_j^1} \times \mathbb{Z}_{n_j^2}), \nu) \rtimes \bigoplus_{j=0}^{\infty} (\mathbb{Z}_{n_j^1} \oplus \mathbb{Z}_{n_j^2})$ , where the action is given by the natural addition, and the measure  $\nu$  is the product measure of  $\nu_j^1$  on  $\mathbb{Z}_{n_j^1}$  and  $\nu_j^2$  on  $\mathbb{Z}_{n_j^2}$ .  $\nu_j^1(pt) = 1/n_j^1$ ,  $\nu_j^2(pt) = 1/n_j^2$ . Under this isomorphism,  $\text{Ad}(\langle t, h' \rangle)\alpha_t$  is conjugate to

$$\bigotimes_{j=0}^{\infty} (\text{Ad}(\text{diag}(\langle t, \rho_{j,0}^1 \rangle, \dots, \langle t, \rho_{j,n_j^1}^1 \rangle)) \otimes \text{Ad}(\text{diag}(\langle t, \rho_{j,0}^2 \rangle, \dots, \langle t, \rho_{j,n_j^2}^2 \rangle))),$$

by (10) and (11). Setting

$$\alpha'_t = \bigotimes_{j=0}^{\infty} \text{Ad}(\text{diag}(\langle t, \rho_{j,0}^2 \rangle, \dots, \langle t, \rho_{j,n_j^2}^2 \rangle)),$$

we know by (12) that  $\alpha$  is cocycle conjugate to  $\tilde{\beta} \otimes \alpha'$ , which is of the infinite tensor product type. We also know  $\alpha \otimes \beta$  is cocycle conjugate to  $\beta \otimes \tilde{\beta} \otimes \alpha' \cong \tilde{\beta} \otimes \alpha'$ , which is cocycle conjugate to  $\alpha$ . Q.E.D.

**COROLLARY 1.9.** *If an action  $\alpha$  of a separable locally compact abelian group  $G$  on the hyperfinite type II<sub>1</sub> factor  $\mathcal{R}$  fixes a Cartan subalgebra and  $\Gamma(\alpha) = \hat{G}$ , then this  $\alpha$  is unique up to cocycle conjugacy.*

*Proof.* Suppose  $\alpha, \beta$  be actions as in the statement. We may assume these are of the above type, and by Theorem 1.6, we may also assume these are of the infinite tensor product type by changing these within their cocycle conjugacy classes if necessary. Now again by Theorem 1.6, both  $\alpha$  and  $\beta$  are cocycle conjugate to  $\alpha \otimes \beta$ , thus  $\alpha$  and  $\beta$  are cocycle conjugate. Q.E.D.

We can apply the above technique to the hyperfinite type II<sub>∞</sub> factor  $\mathcal{R}_{0,1}$ , too.

**PROPOSITION 1.10.** *If  $\alpha$  is an action of a separable locally compact abelian group  $G$  on the hyperfinite type II<sub>∞</sub> factor  $\mathcal{R}_{0,1}$  which fixes a Cartan subalgebra and  $\Gamma(\alpha) = \hat{G}$ , then this  $\alpha$  is unique up to cocycle conjugacy.*

*Proof.* We may assume  $\mathcal{R}_{0,1} = L^\infty(X, \mu) \rtimes_\sigma \mathbb{Z}$ , where  $T$  is a measure preserving transformation on a measure space  $L^\infty(X, \mu)$ ,  $\mu(X) = \infty$ , and  $\alpha$  is given by

$$\begin{cases} \alpha_t(\varphi) = \varphi, & \text{for } \varphi \in L^\infty(X, \mu) \\ \alpha_t(u) = \langle t, h(x) \rangle u, \end{cases}$$

Then choose a sequence of mutually disjoint measurable sets  $\{X_n\}_{n \in \mathbb{N}}$  in  $X$  such that  $X = \bigcup_{n \in \mathbb{N}} X_n$ ,  $\mu(X_n) = 1$ . Choose an open neighborhood  $W \subseteq \hat{G}$  of 0. Then by applying Lemma 1.5, we get  $S_n \in [T]$  such that  $S_0 = \text{id}$ ,  $S_n(X_0) = X_n$ , and  $F(h; x, S_n) \in W$  for almost all  $x \in X_0$ . Define  $h_1(x) = F(h; S_n^{-1}x, x)$  for  $x \in X_n$ , and consider the induced transformation  $T_0$  on  $X_0$ , the reduced cocycle  $c_{X_0}$ , and the action  $\alpha'$  given by  $c_{X_0}$  on the hyperfinite type II<sub>1</sub> factor  $L^\infty(X_0, \mu) \rtimes_{T_0} \mathbb{Z}$  as above. Then  $\text{Ad}(\langle t, h_1 \rangle)\alpha$  is given by  $\alpha' \otimes i$  and  $\Gamma(\alpha') = \hat{G}$ , where  $i$  stands for the trivial action on the type I<sub>∞</sub> factor. If  $\beta$  is another action as in the proposition, we get  $\beta'$  similarly. Now  $\alpha'$  and  $\beta'$  are cocycle conjugate, so we know that  $\alpha$  and  $\beta$  are cocycle conjugate. Q.E.D.

## 2. ALMOST PERIODIC PRIME ACTION OF LOCALLY COMPACT ABELIAN GROUPS WITH $(\mathcal{R}^\alpha)' \cap \mathcal{R} = \mathbf{CI}$

As an application of the result in Section 1, we consider almost periodic prime actions of separable locally compact abelian groups with  $(\mathcal{R}^\alpha)' \cap \mathcal{R} = \mathbf{CI}$ . We keep denoting a separable locally compact abelian group by  $G$ , and consider an action  $\alpha$  of  $G$  on the hyperfinite type II<sub>1</sub> factor  $\mathcal{R}$ . We say  $\alpha$  is *prime* if the fixed point algebra  $\mathcal{R}^\alpha$  is a factor. We define an eigenspace  $\mathcal{R}(p)$  for  $p \in \hat{G}$  by

$$\mathcal{R}(p) = \{x \in \mathcal{R} \mid \alpha_g(x) = \langle g, p \rangle x, \text{ for all } g \in G\},$$

and the pure point spectrum  $\text{Sp}_d(\alpha)$  by

$$\text{Sp}_d(\alpha) = \{p \in \hat{G} \mid \mathcal{R}(p) \neq 0\}.$$

We say  $\alpha$  is almost periodic if the linear span of the subspaces  $\mathcal{R}(p)$ ,  $p \in \hat{G}$ , is weakly dense in  $\mathcal{R}$ . (Definition 7.3 in [16] and Definition 7.1 in [25].) Note the assumption of the existence of a faithful normal  $\alpha$ -invariant state in definitions of [16] and [25] is unnecessary here because we consider the type II<sub>1</sub> factor. In this section, we assume  $\alpha$  is a faithful, almost periodic action of  $G$  on the hyperfinite type II<sub>1</sub> factor  $\mathcal{R}$  with  $(\mathcal{R}^\alpha)' \cap \mathcal{R} = \mathbf{CI}$ , and we will show the uniqueness up to cocycle conjugacy of this type of action. (In this case,  $\alpha$  is prime by Theorem of [18].) We need the following lemma.

LEMMA 2.1. Let  $H$  be a separable compact abelian group, and let  $\hat{H} = \{\lambda_n \mid n \in \mathbf{N}\}$ . Define the infinite tensor product type action  $\sigma$  of  $H$  on the hyperfinite type  $\text{II}_1$  factor  $\mathcal{R}$  by

$$\sigma_h = \bigotimes_{j, n \in \mathbf{N}} \text{Ad} \begin{pmatrix} 1 & 0 \\ 0 & \langle h, \lambda_{n,j} \rangle \end{pmatrix},$$

where  $\lambda_{n,j} = \lambda_n$  for all  $j \in \mathbf{N}$ . Then this is faithful, the Connes spectrum of  $\sigma$  is equal to  $\hat{H}$ , and  $(\mathcal{R}^\sigma)' \cap \mathcal{R} = \mathbb{C}I$ .

*Proof.* It is trivial that this  $\sigma$  is faithful. First we calculate the Connes spectrum  $\Gamma(\sigma)$ . Although  $\Gamma(\sigma) = \hat{H}$  follows from  $(\mathcal{R}^\sigma)' \cap \mathcal{R} = \mathbb{C}I$  by Theorem of [19], we will need this type of calculation later. We make a sequence  $\{\mu_n\}_{n \in \mathbf{N}}$  by renumbering the double sequence  $\{\lambda_{n,j}\}_{n,j \in \mathbf{N}}$ . Set  $X = \prod_{n=1}^{\infty} \{0, 1\}$ , and let  $\mu$  be the product measure of the measure  $\nu$  on  $\{0, 1\}$ ,  $\nu(\{0\}) = \nu(\{1\}) = 1/2$ . We define an equivalence relation  $x \sim y$  for  $x = (x_n), y = (y_n) \in X$  by

$$x \sim y \iff x_n = y_n \quad \text{for all sufficiently large } n's.$$

Then this induces a groupoid, and we can define an  $\hat{H}$ -valued 1-cocycle  $c$  by  $c(x, y) = \sum_{n=1}^{\infty} (y_n - x_n) \mu_n$  for  $x \sim y \in X$ . (Note that the sum is actually a finite sum.) Because this groupoid is amenable, this cocycle is of the type considered in Section 1, and the obtained action is exactly  $\sigma$ . Thus it is enough to show  $r^*(c) = \hat{H}$  by Proposition 1.2, and it is also enough to show that for every  $E \subseteq X$  there exists an integer  $k$  such that if  $n > k$ , then  $\mu_n \in r^*(c_E)$ . (Here  $c_E$  is the restriction of the above cocycle  $c$  to  $E$ . See Proposition 7.6 in [6].)

Fix  $E \subseteq X$ ,  $\mu(E) \neq 0$  and take  $\varepsilon = \mu(E)/9$ . Then there exist an integer  $k$  and  $F \subseteq \prod_{j=1}^k \{0, 1\} \subseteq X$  such that  $\mu(E \Delta F) < \varepsilon$ . (We identify a set  $A \subseteq \prod_{j=1}^k \{0, 1\}$  with  $A \times \prod_{j=k+1}^{\infty} \{0, 1\}$ .) Let  $F = \prod_{l=1}^L F_l$ , where each  $F_l$  is a singleton in  $\prod_{j=1}^k \{0, 1\}$ . Note that each  $F_l$  has measure  $1/2^k$ . We show  $\mu(g_n(E \cap G_n^0) \cap E \cap G_n^1) \neq 0$  for  $n > k$ , where  $g_n = (0, \dots, 0, 1, 0, \dots) \in \bigoplus_{j=1}^{\infty} \mathbb{Z}_2$  (1 is at the  $n$ -th entry),  $G_n^0 = \prod_{j=1}^{n-1} \{0, 1\} \times \{0\} \times \prod_{j=n+1}^{\infty} \{0, 1\}$ , and  $G_n^1 = \prod_{j=1}^{n-1} \{0, 1\} \times \{1\} \times \prod_{j=n+1}^{\infty} \{0, 1\}$ . Suppose it was zero. Setting  $E_l = E \cap F_l$ , we get  $\mu(g_n(E_l \cap G_n^0) \cap E_l \cap G_n^1) \neq 0$  for each  $l$ . It

implies  $\mu(E_l) \leq 1/2^{k+1}$ . Thus  $\mu(F_l - E_l) \geq 1/2^{k+1}$ , and we get

$$\mu(E \Delta F) \geq \sum_{l=1}^L \mu(F_l - E_l) \geq \sum_{l=1}^L 1/2^{k+1} = \mu(F)/2,$$

which implies  $(\sqrt{2} + 1)\mu(E)^{1/2}/3 \geq \mu(E)^{1/2}$ , but this is a contradiction. Thus  $\mu(g_n^{-1}(E \cap G_n^0) \cap E \cap G_n^1) \neq 0$ . Because  $c(x, g_n x) = \mu_n$  for  $x \in E \cap G_n^0 \cap g_n^{-1}(E \cap G_n^1)$ , we get  $\mu_n \in r^*(c_E)$ . Thus we now have  $\Gamma(\sigma) = \hat{H}$ .

It is known that the dual action of the free action of  $\mathbb{Z}$  on  $\mathcal{R}$  is conjugate to the infinite tensor product type action  $\bigotimes_{j=1}^{\infty} \text{Ad} \begin{pmatrix} 1 & 0 \\ 0 & z \end{pmatrix}$ ,  $z \in \mathbb{T}$ , of  $\mathbb{T}$ . Because each element of  $\hat{G}$  appears for infinitely many times in  $\{\mu_n\}$ , we can write  $\mathcal{R} \cong \bigotimes_{j=1}^{\infty} \mathcal{R}_j, \bigotimes_{j=1}^{\infty} \mathcal{P}_j \subseteq \mathcal{R}^\sigma$ , where  $\mathcal{R}_j \cong \mathcal{P}_j \cong \mathcal{R}, \mathcal{P}_j \cap \mathcal{R}_j = \text{CI}$  for all  $j$ . This implies  $(\mathcal{R}^\sigma)' \cap \mathcal{R} = \text{CI}$ . Q.E.D.

Now we can prove the following theorem.

**THEOREM 2.2.** *For a separable locally compact abelian group  $G$ , a faithful almost periodic action  $\alpha$  with  $(\mathcal{R}^\alpha)' \cap \mathcal{R} = \text{CI}$  is unique up to cocycle conjugacy.*

*Proof.* By Proposition 7.3 in [25],  $\text{Sp}_d(\alpha)$  is a countable dense subgroup of  $\hat{G}$ , and if we set  $H = \text{Sp}_d(\alpha)^\wedge$ , there exists an action  $\beta$  of the separable compact abelian group  $H$  on  $\mathcal{R}$  such that  $\alpha_g = \beta_{\iota(g)}$ , where  $\iota$  is a natural dense embedding  $G \subseteq H$ . Because  $\mathcal{R}^\alpha$  is a factor by assumption and  $\iota$  is a dense embedding,  $\mathcal{R}^\beta = \mathcal{R}^\alpha$  is also a factor. By Theorem 5.2 in [25], we know that this  $\beta$  is conjugate to the action  $\sigma$  in Lemma 2.1 because the fixed point algebras of the both have trivial relative commutants. By  $\alpha_g = \beta_{\iota(g)}$ ,  $\alpha$  is also of infinite tensor product type. Because  $\Gamma(\alpha) = \hat{G}$  by the almost periodicity and primeness, we know that this  $\alpha$  is unique up to cocycle conjugacy by Corollary 1.9. Q.E.D.

**REMARK 2.3.** A faithful almost periodic ergodic action is a particular case of prime action. A classification of this type of actions up to conjugacy was given by Theorem 7.4 in [16], and this result was extended to faithful almost periodic prime actions by Theorem 7.4 in [25]. Their invariants are  $\text{Sp}_d(\alpha)$  and a symplectic bicharacter  $\chi_\alpha$  in [16], and a fixed point algebra  $\mathcal{M}^\alpha$  and  $N(\alpha) = \text{Sp}_d(\alpha | (\mathcal{M}^\alpha)' \cap \mathcal{M})$  in addition to these two in [25]. Theorem 7.5 in [25] says there is only one action of the type of Theorem 2.2 up to conjugacy for each  $\text{Sp}_d(\alpha)$ . Our Theorem 2.2 shows that the cocycle conjugacy class of this type of action does not depend on the embedding of  $G$  into  $H = \text{Sp}_d(\alpha)^\wedge$ , and these actions are unique if we consider cocycle conjugacy.

**REMARK 2.4.** If we assume only the almost periodicity in Theorem 2.2, the conclusion is valid. Suppose  $u$  and  $v$  are generating unitaries of the hyperfinite type

$\text{II}_1$  factor  $\mathcal{R}$  with the relation  $uv = e^{2\pi i\theta}vu$ , where  $\theta$  is an irrational number. A one-parameter automorphism group  $\alpha$  is defined by  $\alpha_t(u) = e^{2\pi it}u$ ,  $\alpha_t(v) = e^{2\pi i\theta t}v$ . It is easy to see that this is faithful and almost periodic. Because  $\alpha_1(u) = u$  and  $\alpha_1(v) = e^{2\pi i\theta}v$ , we get  $\alpha_1 = \text{Ad}(u)$ , which is inner. But it can be shown for every almost periodic one-parameter automorphism group  $\beta$  of  $\mathcal{R}$  with  $(\mathcal{R}^\beta)' \cap \mathcal{R} = \mathbb{C}I$ ,  $\beta_t$  is outer for every  $t \neq 0$ . (See Example 4.5.) This implies the above  $\alpha$  cannot be cocycle conjugate to the actions of the type of Theorem 2.2.

### 3. ONE-PARAMETER AUTOMORPHISM GROUPS OF $\mathcal{R}$

In this section we study the case where the group  $G$  in Sections 1 and 2 is the real number group  $\mathbf{R}$ .

In Theorem 0.1 in [13], we classified one-parameter automorphism group  $\alpha$  of the hyperfinite type  $\text{II}_1$  and  $\text{II}_\infty$  factors  $\mathcal{R}$ ,  $\mathcal{R}_{0,1}$  up to stable conjugacy under the assumption  $\Gamma(\alpha) \neq \mathbf{R}$ . Thus we have a complete classification for one-parameter automorphism groups of  $\mathcal{R}$ ,  $\mathcal{R}_{0,1}$  fixing a Cartan subalgebra by this result, Corollary 1.9, and Proposition 1.10. In [13], we considered the ergodic flow on  $\mathcal{Z}(\mathcal{M} \rtimes_\sigma \mathbf{R})$  given by  $\hat{\alpha}$  and the type of  $\mathcal{M} \rtimes_\sigma \mathbf{R}$  as complete invariants for  $\mathcal{M} = \mathcal{R}$ ,  $\mathcal{R}_{0,1}$ . In this section, we show all the ergodic flows occur as this invariant by a similar construction to actions in Section 1. Note that we showed in [13] the type of the crossed product algebra is of  $\text{II}_\infty$  or  $\text{I}_\infty$  unless  $\alpha_t$  is inner for every  $t \in \mathbf{R}$ . Because we have an invariant trace on the crossed product algebra, if the crossed product is of type  $\text{I}_\infty$ , only measure preserving ergodic flows can occur. In the following, we show this is the only restriction on these complete invariants.

**PROPOSITION 3.1.** *In the above context, all the measure preserving ergodic flows occur as  $\hat{\alpha}$  on  $\mathcal{Z}(\mathcal{M} \rtimes_\alpha \mathbf{R})$  if  $\mathcal{M} \rtimes_\alpha \mathbf{R}$  is of type  $\text{I}_\infty$ , and all the ergodic flows occur as  $\hat{\alpha}$  on  $\mathcal{Z}(\mathcal{M} \rtimes_\sigma \mathbf{R})$  if  $\mathcal{M} \rtimes_\alpha \mathbf{R}$  is of type  $\text{II}_\infty$ .*

*Proof.* Suppose an ergodic flow  $T_t$  on a measure space  $(Y, \nu)$  is given. First assume this is measure preserving. Then by Theorem of Ambrose-Kakutani (see [12]), there exist an ergodic measure preserving transformation  $T$  on the measure space  $(Y, \mu)$  and a positive measurable function  $h$  on  $X$  such that  $T_t$  on  $Y$  is conjugate to the flow under the ceiling function  $h$  over the base  $X$ . Construct a one-parameter automorphism group  $\alpha$  for this  $X, T$ , and  $h$  as in Section 1. Then by Proposition 1.3, we know that  $\hat{\alpha}$  on  $\mathcal{Z}(\mathcal{R} \rtimes_\alpha \mathbf{R})$  is conjugate to  $T_t$ . It can be shown as in the argument after Lemma 1.1 in [13] that the crossed product  $\mathcal{R} \rtimes_\alpha \mathbf{R}$  is of type  $\text{I}_\infty$ . If we consider  $\alpha \otimes i$  on  $\mathcal{R} \overline{\otimes} \mathcal{R}$ , where  $i$  is trivial action of  $\mathbf{R}$  on the second copy of  $\mathcal{R}$ , we get a type  $\text{II}_\infty$  crossed product algebra and the same flow as  $\hat{\alpha}$  on  $\mathcal{Z}(\mathcal{M} \rtimes_\alpha \mathbf{R})$ .

Next assume there is no measure on  $Y$  that is equivalent to  $\nu$  and preserved by  $T_t$ . By Theorem of Ambrose-Katukani-Krengel-Kubo (see [12]), there exist an ergodic measurable transformation  $T$  on a measure space  $(X, \mu)$  and a positive measurable function  $h$  on  $X$  such that  $T_t$  on  $Y$  is conjugate to the flow under the ceiling function  $h$  over the base  $X$ . Take and fix an action  $\theta$  of  $\mathbf{R}$  on  $\mathcal{R}_{0,1}$  such that we have  $\text{tr} \circ \theta_t = e^t \text{tr}$  where  $\text{tr}$  is the trace on  $\mathcal{R}_{0,1}$  and  $t \in \mathbf{R}$ . (See [23].) We define an automorphism  $\sigma$  of  $L^\infty(X, \mu) \overline{\otimes} \mathcal{R}_{0,1}$  by  $\sigma(y) = \theta_{-\log m(x)} y$  for  $x \in X$ , where this define a map from  $\mathcal{R}_{0,1}(x)$  to  $\mathcal{R}_{0,1}(Tx)$  in

$$L^\infty(X, \mu) \overline{\otimes} \mathcal{R}_{0,1} = \int_X^\oplus \mathcal{R}_{0,1}(x) dx, \quad \mathcal{R}_{0,1}(x) = \mathcal{R}_{0,1},$$

and  $m(x)$  is the value of Radon-Nikodym derivative of  $T$  at  $x \in X$ . Then this  $\sigma$  is trace preserving on  $L^\infty(X, \mu) \overline{\otimes} \mathcal{R}_{0,1}$ , where the trace is given by  $\mu$  and  $\text{tr}$ . By Lemma 7.11.10 in [20] and the ergodicity of  $T$ , we know that  $(L^\infty(X, \mu) \overline{\otimes} \mathcal{R}_{0,1}) \rtimes_\sigma \mathbf{Z}$  is a factor. Because it has a trace and it is infinite, it is isomorphic to  $\mathcal{R}_{0,1}$ . On this  $(L^\infty(X, \mu) \overline{\otimes} \mathcal{R}_{0,1}) \rtimes_\sigma \mathbf{Z} \simeq \mathcal{R}_{0,1}$ , we can define a one-parameter automorphism  $\alpha_t$  by

$$\begin{cases} \alpha_t(\varphi) = \varphi & \text{for } \varphi \in L^\infty(X, \mu) \overline{\otimes} \mathcal{R}_{0,1} \\ \alpha_t(u) = \langle t, h(x) \rangle u, \end{cases}$$

where  $u$  is the implementing unitary in the crossed product. Then by a similar argument to Proposition 1.3, we know that  $\hat{\alpha}$  on  $\mathcal{Z}(\mathcal{R}_{0,1} \rtimes_\alpha \mathbf{R})$  is conjugate to  $T_t$ .

Q.E.D.

In group actions, the trivial relative commutant property  $\mathcal{R}' \cap \mathcal{R} \rtimes G = \mathbf{C}I$  has been important. We prove this property for one-parameter automorphism groups of the type in Section 1.

**PROPOSITION 3.2.** *If a one-parameter automorphism group  $\alpha$  of the hyperfinite type  $\text{II}_1$  factor  $\mathcal{R}$  fixes a Cartan subalgebra and  $\Gamma(\alpha) = \mathbf{R}$ , then this  $\alpha$  has the trivial relative commutant property,  $\mathcal{R}' \cap \mathcal{R} \rtimes_\alpha \mathbf{R} = \mathbf{C}I$ .*

*Proof.* Because the trivial relative commutant property is invariant under cocycle conjugacy, we may assume  $\alpha$  is of infinite tensor product type by Theorem 1.6. Thus we have an increasing sequence  $\mathcal{M}_n$  of matrix algebras in  $\mathcal{R}$  such that  $\alpha(\mathcal{M}_n) = \mathcal{M}_n$  and  $\bigvee_n \mathcal{M}_n = \mathcal{R}$ . Suppose  $x \in \mathcal{R}' \cap \mathcal{R} \rtimes_\alpha \mathbf{R}$ . Let  $\mathcal{E}_n$  be the conditional expectation from  $\mathcal{R} \rtimes_\alpha \mathbf{R}$  onto  $\mathcal{M}_n \rtimes_\alpha \mathbf{R}$ . Now  $x \in \mathcal{R}' \cap \mathcal{R} \rtimes_\alpha \mathbf{R}$  implies  $\mathcal{E}_n(x) \in \mathcal{M}'_n \cap \mathcal{M}_n \cap \mathcal{M}_n \rtimes_\alpha \mathbf{R} = \lambda(\mathbf{R})$ , where  $\lambda$  denotes the representation of  $\mathbf{R}$  in the crossed product algebra. (Note that  $\alpha|_{\mathcal{M}_n}$  is inner.) Thus  $x = \lim_{n \rightarrow \infty} \mathcal{E}_n(x) \in \lambda(\mathbf{R}) = \lambda(\mathbf{R})'$ , hence we have  $x \in \mathcal{R}' \cap \lambda(\mathbf{R})' = \mathbf{C}I$  because  $\mathcal{R} \rtimes_\alpha \mathbf{R}$  is a factor by  $\Gamma(\alpha) = \mathbf{R}$ .

Q.E.D.

In general, it is not very easy to compute the asymptotic range in Proposition 1.2. But we have the following example for one-parameter automorphism groups.

**EXAMPLE 3.3.** Let  $\theta$  be an irrational number,  $0 < \theta < 1$ , and consider the torus  $\mathbb{T} = [0, 1)$  with the Lebesgue measure, and an ergodic transformation  $T$  on  $X$  defined by  $Tx = x + \theta$ . Take a number  $c$ ,  $0 < c < 1$ ,  $c \notin \mathbb{Q} + \mathbb{Q}\theta$ , and define a function  $h(x) = \chi_{[0,c]}(x) - c$ . Define an action  $\alpha$  on  $\mathcal{R} = L^\infty(\mathbb{T}, \mu) \rtimes \mathbb{Z}$  by  $h$  as above. Then by Theorem A in [17], we have  $r(h, T) = E$ , where  $E$  is the closed subgroup of  $\mathbb{R}$  generated by 1 and  $c$ , which is  $\mathbb{R}$ , and we get an example for  $\Gamma(\alpha) = \mathbb{R}$ . Corollary 1.9 shows that the cocycle conjugacy class of this action does not depend on choice of an irrational  $c$ .

We also have the following for almost periodic prime actions of  $\mathbb{R}$ .

**COROLLARY 3.4.** *For an almost periodic one-parameter automorphism group  $\alpha$  on the hyperfinite type  $\text{II}_1$  factor  $\mathcal{R}$  with  $(\mathcal{R}^\alpha)' \cap \mathcal{R} = \mathbb{C}I$ , we have the trivial relative commutant property  $\mathcal{R}' \cap \mathcal{R} \rtimes_\alpha \mathbb{R} = \mathbb{C}I$ .*

*Proof.* It is clear by Theorem 2.2 and Proposition 3.2.

We consider the next example of an almost periodic one-parameter automorphism group on the hyperfinite type  $\text{II}_1$  factor  $\mathcal{R}$ .

**EXAMPLE 3.5.** Take a free action  $\beta$  of  $\mathbb{Z}^2$  on  $\mathcal{R}$ , and make the crossed product algebra  $\mathcal{R} \rtimes_\beta \mathbb{Z}^2$ , which is isomorphic to  $\mathcal{R}$ . A one-parameter automorphism group  $\alpha$  can be defined by  $\alpha_t(x) = x$  for  $x \in \mathcal{R}$ ,  $\alpha_t(u) = e^{it\mu}u$  and  $\alpha_t(v) = e^{it\lambda}v$ , where  $u$  and  $v$  are the implementing unitaries for  $\mathbb{Z}^2$ , and  $\lambda$  and  $\mu$  are nonzero numbers with  $\lambda/\mu \notin \mathbb{Q}$ . Then it is easy to show this is faithful, almost periodic, and  $(\mathcal{R}^\alpha)' \cap \mathcal{R} = \mathbb{C}I$ . Thus by Theorem 2.2, this is cocycle conjugate to the action in Example 3.3.

#### 4. THE CAR $C^*$ -ALGEBRA AND QUASI-FREE ACTIONS OF $\mathbb{R}$

As an application of the theorem in Section 1, we will classify quasi-free actions arising from the CAR  $C^*$ -algebra.

We introduce a quasi-free action of  $\mathbb{R}$  on the hyperfinite type  $\text{II}_1$  factor  $\mathcal{R}$  as follows.

Take a separable Hilbert space  $\mathcal{H}$ . There exists the Fock representation  $f \mapsto a(f) \in \mathcal{L}(\mathcal{K})$  on another Hilbert space  $\mathcal{K}$ , which satisfies

- (1)  $a(\alpha f + \beta g) = \alpha a(f) + \beta a(g)$ ,  $\alpha, \beta \in \mathbb{C}$ ,
- (2)  $a(f)a(g) + a(g)a(f) = 0$ ,
- (3)  $a(f)^*a(g) + a(g)a(f)^* = (f | g)_\mathcal{H}I_\mathcal{K}$ .



Then  $a(f)$ 's generate a  $C^*$ -algebra, and if  $\{f_n\}_{n \geq 1}$  is a complete orthonormal basis for  $\mathcal{H}$ , we get the following correspondence between this  $C^*$ -algebra and the  $2^\infty$  UHF algebra:

$$\begin{aligned}
 a(f_1) &\longleftrightarrow \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \\
 a(f_2) &\longleftrightarrow \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \otimes \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \\
 a(f_3) &\longleftrightarrow \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \otimes \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \otimes \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \\
 &\vdots
 \end{aligned}$$

We get the hyperfinite type II<sub>1</sub> factor  $\mathcal{R}$  by taking a weak closure with respect to the trace. If we have a one-parameter unitary group  $\{U_t\}_{t \in \mathbb{R}}$  on  $\mathcal{H}$ , then

$$\alpha_t(P(a(f_1), \dots, a(f_n)^*)) = P(a(U_t f_1), \dots, a(U_t f_n)^*),$$

where  $P$  is a (non-commutative) polynomial, defines a one-parameter automorphism group on the CAR  $C^*$ -algebra. This extends to a one-parameter automorphism group on  $\mathcal{R}$ , and we denote it by  $\alpha$ , too. We call this a quasi-free action of  $\mathbb{R}$  on  $\mathcal{R}$ , and classify this type of actions in this section.

In the above context, let  $H$  be a self-adjoint operator such that  $e^{iHt} = U_t$ . Then by von Neumann's theorem (see Theorem X.2.1 in [11]) there exists a Hilbert-Schmidt class self-adjoint operator  $V$  such that a self-adjoint operator  $K = H + V$  has pure point spectrum. Then the one-parameter unitary group  $e^{iKt}$  defines another one-parameter automorphism group  $\beta$  on  $\mathcal{R}$ . We have the following for these two actions. (See also p.315 in [22].)

**THEOREM 4.1.** *Let  $\alpha$  and  $\beta$  be quasi-free actions of  $\mathbb{R}$  on  $\mathcal{R}$  corresponding to  $e^{iHt}$  and  $e^{iKt}$  respectively as above. Then  $\alpha$  is cocycle conjugate to  $\beta$ , and  $\beta$  is of the infinite tensor product type.*

*Proof.* Let  $V = \sum_{n=1}^\infty \lambda_n E_n$ , where  $E_n$  is a mutually orthogonal rank-one projection onto a subspace spanned by  $f_n$ . We may assume  $\{f_n\}_{n \geq 1}$  is a complete orthogonal basis of  $\mathcal{H}$ . We know that  $\lambda_n$ 's are real and  $\sum_{n=1}^\infty \lambda_n^2 < \infty$ . Let  $\alpha_t^{(n)}$  be the one-parameter automorphism group on  $\mathcal{R}$  corresponding to the one-parameter unitary

group  $\exp it \left( H + \sum_{j=1}^n \lambda_j E_j \right)$ . (We use the convention  $\alpha^{(0)} = \alpha$ .) We use the above correspondence between the CAR  $C^*$ -algebra and the infinite tensor product of copies of the  $2 \times 2$  matrix algebra for the orthonormal sequence  $\{f_n\}_{n \geq 1}$ . Define for  $n \geq 0$ ,

$$u_t^{(n)} = \text{Exp}_r \left( \int_0^t \alpha_s^{(n)} \left( 1 \otimes \cdots \otimes 1 \otimes \begin{pmatrix} i\lambda_{n+1}/2 & 0 \\ 0 & -i\lambda_{n+1}/2 \end{pmatrix} \right) ds \right),$$

where  $\text{Exp}_r$  means an expansional. (See §2 in [1].) By Theorem 2 in [1], it is an  $\alpha^{(n)}$ -unitary cocycle. If  $g$  is in  $\text{Dom}(H) = \text{Dom}(K) \subseteq \mathcal{H}$ , then we have

$$\begin{aligned} \delta_{\alpha^{(n+1)}}(a(g)) &= \\ &= a \left( i \left( H + \sum_{j=1}^n \lambda_j E_j + \lambda_{n+1} E_{n+1} \right) g \right) = \\ &= \delta_{\alpha^{(n)}}(a(g)) + \\ &+ \frac{d}{dt} \left( \text{Ad} \left( \exp it \left( 1 \otimes \cdots \otimes 1 \otimes \begin{pmatrix} \lambda_{n+1}/2 & 0 \\ 0 & -\lambda_{n+1}/2 \end{pmatrix} \right) \right) a(g) \right) \Big|_{t=0} = \\ &= \delta_{\alpha^{(n)}}(a(g)) + \left[ 1 \otimes \cdots \otimes 1 \otimes \begin{pmatrix} i\lambda_{n+1}/2 & 0 \\ 0 & -i\lambda_{n+1}/2 \end{pmatrix}, a(g) \right], \end{aligned}$$

where  $\delta$ 's stand for derivations. Thus  $\alpha_t^{(n+1)}(a(g)) = \text{Ad}(u_t^{(n)})\alpha_t^{(n)}(a(g))$  by Theorem 2 in [1], and thus we get  $\alpha_t^{(n+1)} = \text{Ad}(u_t^{(n)})\alpha_t^{(n)}$ . Thus if we set  $v_t^{(n)} = u_t^{(n)} \cdots u_t^{(0)}$ , it is an  $\alpha$ -unitary cocycle. Note that we have

$$v_t^{(n)} = \text{Exp}_r \left( \int_0^t \alpha_s \left( i \begin{pmatrix} \lambda_1/2 & 0 \\ 0 & -\lambda_1/2 \end{pmatrix} \odot \cdots \odot \begin{pmatrix} \lambda_{n+1}/2 & 0 \\ 0 & -\lambda_{n+1}/2 \end{pmatrix} \right) ds \right),$$

where the operator  $\odot$  is defined by

$$\begin{pmatrix} a & 0 \\ 0 & b \end{pmatrix} \odot \begin{pmatrix} c & 0 \\ 0 & d \end{pmatrix} = \begin{pmatrix} a+c & 0 & 0 & 0 \\ 0 & a+d & 0 & 0 \\ 0 & 0 & b+c & 0 \\ 0 & 0 & 0 & b+d \end{pmatrix}.$$

(This operation is defined similarly for higher dimensional matrices.)

Now we claim that there exists an  $\alpha$ -unitary cocycle  $v_t$  such that  $v_t^{(n)} \rightarrow v_t$  as  $n \rightarrow \infty$  uniformly for  $t$  on every compact set in  $\mathbf{R}$ .

For a given  $\varepsilon$ , choose  $N$  such that  $\left(\sum_{n>N} \lambda_n^2\right)^{1/2} < 2\varepsilon$ . We will show that  $\|v_t^{(n)} - v_t^{(m)}\|_2 \leq \varepsilon t$  for  $n > m > N$ . Set

$$A_{m,n} = 0 \oplus \cdots \oplus 0 \oplus \begin{pmatrix} \lambda_{m+2}/2^0 & 0 \\ 0 & -\lambda_{m+2}/2 \end{pmatrix} \oplus \cdots \oplus \begin{pmatrix} \lambda_{n+1}/2 & 0 \\ 0 & -\lambda_{n+1}/2 \end{pmatrix}$$

Then by Theorem 1 in [1]

$$\begin{aligned} \|v_t^{(n)} - v_t^{(m)}\|_2 &= \left\| \text{Exp}_r \left( \int_0^t \alpha_s^{(m+1)}(A_{m,n}) ds \right) - 1 \right\|_2 = \\ &= \left\| \int_0^t \text{Exp}_r \left( \int_0^s \alpha_r^{(m+1)}(A_{m,n}) dr \right) \alpha_s^{m+1}(A_{m,n}) ds \right\|_2 \leq \\ &\leq \int_0^t \|A_{m,n}\|_2 ds = \left( \sum_{j=m+2}^{n+1} \lambda_j^2 \right)^{1/2} t/2 < \varepsilon t. \end{aligned}$$

Thus  $v_t^n$  converges to an  $\alpha$ -unitary cocycle  $v_t$ . Letting  $n \rightarrow \infty$  in

$$\text{Ad}(v_t^{(n)})\alpha_t(a(g)) = \alpha_t^{(n)}(a(g)) = a \left( \exp it \left( H + \sum_{j=1}^n \lambda_j E_j \right) g \right),$$

we get  $\text{Ad}(v_t)\alpha_t = \beta_t$ .

Because  $K$  has a pure point spectrum, we may assume  $K = \sum_{n=1}^{\infty} \mu_n E_n$ , where  $\mu_n$ 's are real numbers,  $F_n$ 's are mutually orthogonal rank-one projections, and  $\sum_{n=1}^{\infty} F_n = I_{\mathcal{H}}$ . Thus  $\beta_t$  is of the form  $\bigoplus_{n=1}^{\infty} \text{Ad} \left( \exp it \begin{pmatrix} \mu_n/2 & 0 \\ 0 & -\mu_n/2 \end{pmatrix} \right)$ . Q.E.D.

Thus we can apply the result in Section 1 to quasi-free actions. In the following, we consider the above  $\alpha$ , and  $\beta$  which arises from  $K = \sum_{n=1}^{\infty} \mu_n F_n$ . Define a groupoid whose unit is  $X = \prod_{j=1}^{\infty} \{0, 1\}$  as in the proof of Lemma 2.1. We also define an  $\mathbf{R}$ -valued

1-cocycle  $c$  by  $c(x, y) = \sum_{j=1}^{\infty} (x_j - y_j) \mu_j$  for  $x \sim y \in X$ . The obtained one-parameter automorphism group as in Section 1 is  $\beta$ , thus we get the following by Corollary 1.9 and Proposition 1.2.

**COROLLARY 4.2.** *A one-parameter automorphism group  $\alpha$  on  $\mathcal{R}$  arising from the CAR  $C^*$ -algebra in the above way is unique up to cocycle conjugacy if  $\Gamma(\alpha) = \mathbb{R}$ .*

**PROPOSITION 4.3.** *In the above context, the Connes spectra  $\Gamma(\alpha)$  and  $\Gamma(\beta)$  are equal to the asymptotic range  $r^*(c)$ .*

We can also show the trivial relative commutant property (Proposition 3.2) for this type of  $\alpha$  if  $\Gamma(\alpha) = \mathbb{R}$ .

For the computation of examples, we prove the following.

**PROPOSITION 4.4.** *For a one-parameter automorphism group  $\alpha$  on  $\mathcal{R}$  arising from one-parameter unitary group  $e^{iHt}$  in the above way, the essential spectrum  $\sigma_e(H)$  is contained in  $\Gamma(\alpha)$ .*

*Proof.* By von Neumann's theorem again, we get  $H_+ + V = K$  as above. Because  $\sigma_e(H) = \sigma_e(K)$ , we consider  $K = \sum_{n=1}^{\infty} \mu_n P_n$  and  $\beta$  instead of  $H$  and  $\alpha$ .

Then we can prove the statement exactly as in the proof of Lemma 2.1. Q.E.D.

In particular, if the spectrum of  $H$  contains a continuous part, it implies  $\Gamma(\alpha) = \mathbb{R}$  for  $\alpha$  coming from  $e^{iHt}$ , and this type of  $\alpha$  is unique up to cocycle conjugacy.

**EXAMPLE 4.5.** Let  $\mathcal{H} = L^2(\mathbb{R})$ , and define a one-parameter unitary group  $U_t$  by  $U_t f(x) = f(x - t)$  for  $f \in L^2(\mathbb{R})$ . Then we can define a one-parameter automorphism group  $\alpha$  on  $\mathcal{R}$ . (See pp.4-5 in [21]. They construct a similar endomorphism semigroup for the CAR  $C^*$ -algebra.) In this context, above  $H$  is  $i \frac{d}{dx}$ , which has spectrum  $\mathbb{R}$ . Thus by Proposition 4.4, we know  $\Gamma(\alpha) = \mathbb{R}$ .

Take another  $\mathcal{H}' = \mathcal{H}_1 \oplus \mathcal{H}_2$ , where  $\mathcal{H}_1$  and  $\mathcal{H}_2$  are infinite dimensional separable Hilbert spaces. We consider  $H = \lambda_1 I_{\mathcal{H}_1} \oplus \lambda_2 I_{\mathcal{H}_2}$ , where  $\lambda_1$  and  $\lambda_2$  are non-zero real numbers, and  $\lambda_1/\lambda_2$  is irrational. Then we can consider one-parameter unitary group  $e^{iH't}$ , and it induces a one-parameter automorphism group  $\alpha'$  of  $\mathcal{R}$  as above. Because  $\sigma_e(H') = \{\lambda_1, \lambda_2, \}$ ,  $\lambda_1/\lambda_2$  is irrational, and  $\Gamma(\alpha')$  is a closed subgroup of  $\mathbb{R}$ , we know  $\Gamma(\alpha') = \mathbb{R}$  again by Proposition 4.4. Note that this  $\alpha'$  is of the form

$$\bigotimes_{j=1}^{\infty} \text{Ad} \left( \exp it \begin{pmatrix} \lambda_1 & 0 \\ 0 & 0 \end{pmatrix} \right) \otimes \bigotimes_{k=1}^{\infty} \text{Ad} \left( \exp it \begin{pmatrix} \lambda_2 & 0 \\ 0 & 0 \end{pmatrix} \right).$$

(From this expression, it is easy to see that  $\alpha'_t$  is outer for any  $t \neq 0$ .)

Then by Corollary 4.2, we know that  $\alpha$  and  $\alpha'$  are cocycle conjugate. We also know that these are cocycle conjugate to the one-parameter automorphism groups in Examples 3.3 and 3.5.

**REMARK 4.6.** The inclusion in Proposition 4.4 is not the best possible. Actually by very similar arguments to Lemma 4.2, Lemma 4.3, Lemma 5.4 in [2], and Théorème 1 in [3] with Corollaries 4.2 and 4.3 here, we get the following characterization. This corresponds to writing the asymptotic range in the original formulation of the asymptotic ratio set of Araki-Woods.)

A real number  $\lambda$  is in  $\Gamma(\alpha)$  if and only if the following conditions are satisfied:

- (1)  $\{I_n\}_{n \in \mathbb{N}}$  is a family of mutually disjoint finite subsets of  $\mathbb{N}$ ,
- (2)  $K_n^1, K_n^2$  are mutually disjoint subsets of  $\left\{ \sum_{j \in F} \mu_j \mid f \subseteq I_n \right\}$ ,
- (3)  $\psi_n$  is a bijective map from  $K_n^1$  onto  $K_n^2$ ,
- (4)  $\sum_{n=1}^{\infty} |K_n^1|/2^{|I_n|} = \infty$ ,
- (5)  $\lim_{n \rightarrow \infty} \max_{\rho \in K_n^1} |\lambda - (\psi_n(\rho) - \rho)| = 0$ .

In (2),  $\mu_j$ 's are eigenvalues of  $K$  as above.

By this criterion, we can show that the following  $H$  has  $\sigma_e(H) = \{0\}$ , but the one-parameter automorphism group it induces has the Connes spectrum  $\mathbb{R}$ . (Thus it is cocycle conjugate to the actions in Example 4.5.)

Let  $\{a_n\}_{n \in \mathbb{N}}$  be the following sequence:

$$1, 1, \underbrace{\frac{1}{2}, \dots, \frac{1}{2}}_{8 \text{ times}}, \dots, \underbrace{\frac{1}{n}, \dots, \frac{1}{n}}_{n2^n \text{ times}}, \dots$$

Define  $a_{n,m} = \frac{1}{m} a_n$ . We define  $H = \sum_{n,m} a_{n,m} E_{n,m}$ , where  $E_{n,m}$ 's are mutually orthogonal rank-one projections in  $\mathcal{H}$ . Then  $H$  is a compact operator, and  $\sigma_e(H) = \{0\}$ , but by the above criterion, we can show  $1/m$  is in  $\Gamma(\alpha)$  for every positive integer  $m$ . Thus we have  $\Gamma(\alpha) = \mathbb{R}$ .

## REFERENCES

1. ARAKI, H., Expansional in Banach algebras, *Ann. Sci. École Norm. Sup.*, **6**(1973), 67-84.
2. ARAKI, H.; WOODS, E. J., A classification of factors, *Publ. Res. Inst. Math. Sci.*, **3**(1968), 51-130.
3. CONNES, A., Calcul des deux invariants d'Araki et Woods par la théorie de Tomita et Takesaki, *C. R. Acad. Sci. Paris*, **274**(1972), 175-177.
4. CONNES, A., Une classification des facteurs de type III, *Ann. Sci. École Norm. Sup.*, **6**(1973), 133-252.
5. CONNES, A.; FELDMAN, J.; WEISS, B., An amenable equivalence relation is generated by a single transformation, *Ergodic Theory Dynamical Systems*, **1** (1981), 431-450.
6. FELDMAN, J.; MOORE, C. C., Ergodic equivalence relations, cohomology and von Neumann algebras. I, *Trans. Amer. Math. Soc.*, **234**(1977), 289-324.
7. FELDMAN, J.; MOORE, C. C., Ergodic equivalence relations, cohomology and von Neumann algebras. II, *Trans. Amer. Math. Soc.*, **234**(1977), 325-359.
8. HAAGERUP, U., Connes' bicentralizer problem and uniqueness of the injective factor of type III<sub>1</sub>, *Acta Math.*, **158**(1987), 95-147.
9. JONES, V., Actions of finite groups on the hyperfinite type II<sub>1</sub> factor, *Mem. Amer. Math. Soc.*, **237**(1980).
10. JONES, V.; TAKESAKI, M., Actions of compact abelian groups on semifinite injective factors, *Acta Math.*, **153**(1984), 213-258.
11. KATO, T., *Perturbation theory for linear operators*, Springer, 1966.
12. KATOK, A. B., Monotone equivalence in ergodic theory, *Math. USSR-Izv.*, **11**(1977), 99-146.
13. KAWAHIGASHI, Y., Centrally ergodic one-parameter automorphism groups on semifinite injective von Neumann algebras, *Math. Scand.*, to appear.
14. KRIEGER, W., On the Araki-Woods asymptotic ratio set and non-singular transformations of a measure space, Springer Lecture Notes in Math., No. 160, 1970, 158-177.
15. OCNEANU, A., *Actions of discrete amenable groups on factors*, Springer Lecture Notes in Math., No. 1138, 1985.
16. OLESEN, D.; PEDERSEN, G. K.; TAKESAKI, M., Ergodic actions of compact abelian groups, *J. Operator Theory*, **3**(1980), 237-269.
17. OREN, I., Ergodicity of cylinder flows arising from irregularities of distribution, *Israel J. Math.*, **44**(1983), 127-138.
18. PASCHKE, W., Inner product modules arising from compact automorphism groups on von Neumann algebras, *Trans. Amer. Math. Soc.*, **224**(1976), 87-102.
19. PASCHKE, W., Relative commutant of a von Neumann algebra in its crossed product by a group action, *Math. Z.*, **163**(1978), 5-13.
20. PEDERSEN, G. K., *C\*-algebras and their automorphism groups*, London Mathematical Society Monographs, Vol. 14, Academic Press, London, 1979.
21. POWERS, R. T.; ROBINSON, D. W., An index for continuous semigroups of \*-endomorphisms of  $B(H)$ , preprint.
22. SAKAI, S., Developments in the theory of unbounded derivations in C\*-algebras, in *Operator algebras and applications*, Part 2 (Kingston, Ont., 1980), Proc. Sympos. Pure Math., Vol. 38, Amer. Math. Soc., Providence, R. I., 1982, pp. 309-331.
23. TAKESAKI, M., Duality for crossed products and the structure of von Neumann algebras of type III, *Acta Math.*, **131**(1973), 249-310.

24. TAKESAKI, M., *Structure of operator algebras* (Japanese), Iwanami, Tokyo, 1983.
25. THOMSEN, K., *Compact abelian prime actions on von Neumann algebras*, preprint.

YASUYUKI KAWAHIGASHI  
*Department of Mathematics,  
University of California,  
Los Angeles, CA 90024,  
U.S.A.*

Received July 25, 1988.